



INDOXACARB	GROUP	22A	INSECTICIDE
------------	-------	-----	-------------

EPA Reg. No. 279-9629

SUPPLEMENTAL LABELING

FOR FOLIAR APPLICATIONS TO TREE NUTS TO CONTROL CODLING MOTH, NAVEL ORANGEWORM AND PEACH TWIG BORER

THIS SUPPLEMENTAL LABEL EXPIRES 11/30/2023 AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read the label affixed to the container for Avaunt eVo insect control before applying.

Use of Avaunt eVo insect control according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Avaunt eVo insect control.

IMPORTANT

BEFORE USING AVAUNT eVo insect control, READ AND FOLLOW ALL APPLICABLE DIRECTIONS; RESTRICTIONS; AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which does not appear on the package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

Sold By



**FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104**

Crops	Insects	AVAUNT eVo insect control Rate Per Acre		Last Application (Days to Harvest)	REI
		lb ai	Ounces		
Tree Nuts Including: Almond; Pistachio; Walnut, black; Walnut, English cultivars, varieties, and/or hybrids of these	Codling moth* Navel orangeworm** Peach twig borer***	0.075 - 0.11	4.0 - 6.0	5	12 hrs.
<p>TREE NUTS RESTRICTIONS Make no more than 3 applications per season. Do not apply more than 18 oz/A of AVAUNT eVo insect control or 0.33 lb ai/A of indoxacarb-containing products per calendar year. The minimum interval between treatments is 7 days.</p> <p>Note: Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage. Where higher spray volumes are used, apply a higher rate in the specified rate range. Ground Application: Apply in a minimum of 50 gal/A of water by conventional ground equipment. For best results, apply 100-150 gal/A of water. Aerial Application: Apply in a minimum of 10 gal/A of water. Applying by air can result in reduced canopy coverage, which may also reduce efficacy. For best results, use an adjuvant to help increase coverage, penetration and thus performance.</p> <p>*Codling moth – (Walnut) Make initial application at or before peak egg lay for targeted generation. Depending on level of infestation reapply 14 days later as needed. Use higher rates and ground application equipment to achieve thorough coverage.</p> <p>**Navel orangeworm – For NOW, use at May or hull split timing. If targeting a hull split application, apply at 1-5% hull split. Make a second application approximately 10 – 14 days later. Depending on level of pest infestation, use of higher rates in the labeled rate range and multiple applications may be needed.</p> <p>***Peach twig borer - For spring application to overwintering generation: Make application at late dormant (just prior to bud break) to early bloom. For "May spray" applications to the summer generation: Make applications at peak moth flight (timed at or before peak egg lay). Higher rates in the labeled rate range may be needed for high infestations levels and large, dense foliage trees.</p>					

(Derived from AVAUNT eVo insect control Label SL-4187A 010821 11-23-20)

R-4479A 020321 11-23-20

FMC Corporation, 2929 Walnut Street, Philadelphia, PA 19104
 © 2000-2020 FMC Corporation. All rights reserved.