

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

BIFENTHRIN GROUP 3A INSECTICIDE

Supplemental Labeling

This supplemental label expires on December 31, 2024 and must not be used or distributed after this date.



For use on Peach, Pome Fruit and Pomegranate

EPA Reg. No. 279-3108

Active Ingredient:	By Wt.
Bifenthrin: *	10.0%
Other Ingredients:	90.0%
	100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.
ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for BRIGADE WSB Insecticide/Miticide before applying. Use of BRIGADE WSB Insecticide/Miticide according to this supplemental labeling is subject to the use precautions and limitations imposed by the label affixed to the container of BRIGADE WSB Insecticide/Miticide. Carefully follow all precautionary statements and application use directions.



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

PEACH Subgroup 12-12B

Peaches, Nectarines

Pests Controlled	Dosage* oz product/A (lb ai/A)	Directions
Aphids Codling Moth Cutworms Green Fruitworm Leafhoppers Leafminers Leafrollers Lygus spp. Plant Bugs Plum Curculio San Jose Scale (Crawlers) Stink Bugs Tarnished Plant Bugs	6.4 - 32 (0.04 - 0.2)	<p>Application by ground: Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage.</p> <p>Application by air: Apply the specified dosage in a minimum of 10 gallons of finished spray per acre by air.</p> <p>*1 bag = 0.05 lb ai = 8 oz formulated product</p>
Yellow Spider Mite	9.6 - 32 (0.06 - 0.2)	
European Red Mite Twospotted Spider Mite	12.8 - 32 (0.08 - 0.2)	
<p>Restrictions</p> <ul style="list-style-type: none"> • Do not exceed 4 bags/A (0.2 lb ai/A or 32 oz product/A) per application. • Do not exceed 10 bags/A (0.5 lb ai/A or 80 oz product/A) per year with no more than 0.45 lb ai/A applied after petal fall. • Do not make more than 3 applications per year. • Minimum re-treatment interval (RTI) is 30 days. • Do not apply within 14 days of harvest. • Do not graze livestock in treated orchards or cut treated cover crops for feed. 		

POME FRUIT Group 11-10 (except Mayhaw)

Pears, Apples, Azarole, Crabapple, Loquat, Medlar, Quince, Tejocote

Pests Controlled	Dosage* oz product/A (lb ai/A)	Directions
Aphids Codling Moth Cutworms Green Fruitworm Leafhoppers Leafminers Leafrollers Lygus spp. Plant Bugs Plum Curculio San Jose Scale (Crawlers) Stink Bugs Tarnished Plant Bugs	6.4 - 32 (0.04 - 0.2)	<p>Application by ground: Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage.</p> <p>Application by air: Apply the specified dosage in a minimum of 10 gallons of finished spray per acre by air.</p> <p>*1 bag = 0.05 lb ai = 8 oz formulated product</p>
Yellow Spider Mite	9.6 - 32 (0.06 - 0.2)	
European Red Mite Twospotted Spider Mite	12.8 - 32 (0.08 - 0.2)	
<p>Restrictions</p> <ul style="list-style-type: none"> • Do not exceed 4 bags/A (0.2 lb ai/A or 32 oz product/A) per application. • Do not exceed 10 bags/A (0.5 lb ai/A or 80 oz product/A) per year with no more than 0.45 lb ai/A applied after petal fall. • Do not make more than 3 applications per year. • Minimum re-treatment interval (RTI) is 30 days. • Do not apply within 14 days of harvest. • Do not graze livestock in treated orchards or cut treated cover crops for feed. 		

POMEGRANATE

Pests Controlled	Dosage* oz product/A (lb ai/A)	Directions
Katydid navel orangeworm omnivorous leafrollers leaf footed plant bugs fuller rose beetles aphids white scales ground beetles brown marmorated stink bugs	16 – 32 (0.1 – 0.2)	Apply foliar treatments in at least 50 gallons of finished spray per acre. *1 bag = 0.05 lb ai = 8 oz formulated product
Restrictions <ul style="list-style-type: none"> • Do not exceed 4 bags/A (0.2 lb ai/A or 32 oz product/A) per application. • Do not exceed 10 bags/A (0.5 lb ai/A or 80 oz/A) per year. • Do not make more than 3 applications per year. • Do not make applications less than 14 days apart. • Do not apply within 14 days of harvest. 		

FMC and BRIGADE are trademarks of FMC Corporation.
 © 2022 FMC Corporation. All rights reserved.

R-4650 042622 04-15-22