

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

 <p>EPA REG. NO. 279-3108</p>	<u>2(ee) Recommendation</u>
	FOR DISTRIBUTION AND USE ONLY IN New York
	IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING
	THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

For control of spotted lanternfly in listed crops.

CROP	RATE OF APPLICATION*	RESTRICTIONS
Grapes	8 – 16 oz/A (0.05 – 0.10 lb ai/A)	<ul style="list-style-type: none"> Do not apply more than 16 oz of product (0.10 lb ai) per acre per application. Do not apply more than 16 oz of product (0.10 lb ai) per acre per season. Do not make applications less than 7 days apart. Do not apply within 30 days of harvest.

Use Directions:

Spray in sufficient water to provide thorough coverage for best control.

Application by ground: Apply in a minimum of 25 gallons of finished spray per acre.

Application by air: Apply in a minimum of 10 gallons of finished spray per acre.

*1 bag = 0.05 lb ai = 8 oz formulated product.

CROP	RATE OF APPLICATION*	RESTRICTIONS
Hops	8 – 16 oz/A (0.05 – 0.10 lb ai/A)	<ul style="list-style-type: none"> Do not apply more than 16 oz of product (0.10 lb ai) per acre per application. Do not apply more than 48 oz of product (0.30 lb ai) per acre per season. Do not make applications less than 21 days apart. Do not apply within 14 days of harvest.

Use Directions:

Application by ground: For best control, full coverage is essential. Early season



recommendation is 100-150 gallons of finished spray per acre. Late season recommendation is 200-250 gallons of finished spray per acre.

*1 bag = 0.05 lb ai = 8 oz formulated product.

CROP	RATE OF APPLICATION*	RESTRICTIONS
Pears	6.4 – 32.0 oz/A (0.04 – 0.20 lb ai/A)	<ul style="list-style-type: none"> • Do not apply more than 32 oz of product (0.20 lb ai) per acre per application. • Do not apply more than 80 oz of product (0.5 lb ai) per acre per season, with no more than 72 oz of product (0.45 lb ai) per acre applied after petal fall. • Do not graze livestock in treated orchards or cut treated cover crops for feed. • Do not make applications less than 30 days apart. • Do not apply within 14 days of harvest.

Use Directions:

Application by ground: Spray in sufficient water to provide thorough coverage. Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre).

Application by air: Apply in a minimum of 10 gallons of finished spray per acre.

*1 bag = 0.05 lb ai = 8 oz formulated product.

REMARKS:

Thorough coverage is essential to achieve control.

Multiple applications may be required to maintain control and prevent reinfestation from migrating adults into the crop. Spotted lanternflies are mobile and may reinfest crops quickly. Spotted lanternflies should be targeted while in the nymphal stage. If another treatment is needed before the minimum retreatment interval, use another insecticide to maintain control.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED BRIGADE® WSB INSECTICIDE/MITICIDE (EPA REG NO. 279-3108) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

Brigade® and FMC are trademarks of FMC Corporation or an affiliate. ©2018 FMC Corporation.

DR-4104 101718

Expires: 06-30-23



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104