

## Supplemental Label

# Cevya<sup>®</sup>

## Fungicide

### For disease control in bulb vegetables, cucurbit vegetables, fruiting vegetables, and root vegetables (except sugar beet)

This supplemental label expires October 30, 2023, and must not be used or distributed after this date.

#### Active Ingredient\*:

mefentrifluconazole: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazole-1-yl)propan-2-ol .....	34.93%
Other Ingredients: .....	65.07%
Total: .....	100.00%

\* Cevya<sup>®</sup> fungicide contains 3.34 lbs mefentrifluconazole per gallon.

EPA Reg. No. 7969-407

CAUTION/PRECAUCION

#### Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Cevya<sup>®</sup> fungicide** container label, EPA Reg. No. 7969-407, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Cevya** before applying.
- Use of **Cevya** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Cevya**.

#### Application Instructions

Refer to the **Cevya** container label for additional instructions and restrictions.

## Crop Rotation Restrictions

**Rotational Crops:** Please see the following table for crop rotational restrictions.

Crop or Crop Group	Rotation Interval
Brassicas including broccoli, cauliflower and head cabbages (crop group 5-16) Bulb vegetables including bulb onion, spring onion and garlic (crop group 3-07) Cereals, including wheat, oats, barley, triticale, rye, rice and corn (crop group 15 and crop group 16) Cucurbits including cucumber, squash and melons (crop group 9) Fruiting vegetables including tomato, eggplant, peppers (crop group 8-10) Fresh herbs Grass and non-grass animal feeds including alfalfa and clover (crop group 17 and crop group 18) Leafy vegetables including lettuces, spinach and leafy cabbages (crop group 4-16) Legume vegetables including soybean (crop group 6 and crop group 7) Low-growing berries including strawberries (crop group 13-07G) Oilseeds including cotton, sunflower and canola (crop group 20) Peanut Root and tuber vegetables including carrot, potato, beets and sugar beet (crop group 1 and crop group 2) Stalk, stem and leaf petiole vegetables including celery and asparagus (crop group 22) Sugarcane Any other crop labeled for direct application of a product containing mefenftrifluconazole	May be planted immediately following the last application
Other food and feed crops, not listed above	May not be planted in rotation

## Detailed Use Directions

### Labeled Crops

Foliar

- When conditions favor disease or disease pressure is high:
  - Use the shorter specified interval.
  - For a rate range, use the higher specified rate.
- **DO NOT** make more than two (2) sequential applications of **Cevya® fungicide** before alternating to a labeled **non-Group 3** fungicide.

#### Use Rate Conversion

fl ozs product/A	lb mefentrifluconazole/A
3	0.08
4	0.10
5	0.13

**State-specific Restrictions** - Not for sale, distribution, or use in **Nassau and Suffolk counties in New York State.**

## Bulb Vegetables

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Purple blotch and leaf blight <i>Alternaria porri</i>	3 to 5	15	7
	<ul style="list-style-type: none"> <li>• <b>Spray Interval</b> - Apply before the onset of disease and on a minimum interval of 7 days.</li> </ul>			

### Use Restrictions

- **DO NOT** apply more than 5 fl ozs (0.13 lb mefentrifluconazole) per acre per application.
- **DO NOT** make more than 3 applications at 5 fl ozs per acre per year.
- **DO NOT** apply more than 15 fl ozs (0.39 lb mefentrifluconazole) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products.

**Detailed Crop (Group 3-07) List** - chive, fresh leaves; chive, Chinese, fresh leaves; daylily, bulb; elegans hosta; fritillaria, bulb; fritillaria, leaves; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; kurrat; lady's leek; leek; leek, wild; lily, bulb; onion, Beltsville bunching; onion, bulb; onion, Chinese, bulb; onion, fresh; onion, green; onion, macrostem; onion, pearl; onion, potato, bulb; onion, tree, tops; onion, Welsh, tops; shallot, bulb; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.

## Cucurbit Vegetables

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Gummy stem blight <i>Didymella bryoniae</i>	3 to 5	15	0
	Powdery mildew <i>Erysiphe</i> spp., <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp.			
<ul style="list-style-type: none"> <li>• <b>Spray Interval</b> - Apply before onset of disease and on a minimum interval of 7 days.</li> </ul>				

### Use Restrictions

- **DO NOT** apply more than 5 fl ozs (0.13 lb mefentrifluconazole) per acre per application.
- **DO NOT** make more than 3 applications at 5 fl ozs per acre per year.
- **DO NOT** apply more than 15 fl ozs (0.39 lb mefentrifluconazole) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products.

**Detailed Crop (Group 9) List** - chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes true canteloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon.

## Fruiting Vegetables

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Anthracnose <i>Colletotrichum coccodes</i>	3 to 5	15	0
	Black mold <i>Alternaria alternata</i>			
	Early blight <i>Alternaria solani</i>			
	Powdery mildew <i>Leveillula taurica, Oidiopsis taurica</i>			
<p>• <b>Spray Interval</b> - Apply before onset of disease and on a minimum interval of 7 days.</p>				

### Use Restrictions

- **DO NOT** apply more than 5 fl ozs (0.13 lb mefentrifluconazole) per acre per application.
- **DO NOT** make more than 3 applications at 5 fl ozs per acre per year.
- **DO NOT** apply more than 15 fl ozs (0.39 lb mefentrifluconazole) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products.

**Detailed Fruiting Vegetables (crop group 8-10) Crop List** - African eggplant; bush tomato; bell pepper; cocona; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; naranjilla; okra; pea eggplant; pepino; non-bell pepper; roselle; scarlet eggplant; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these.

## Root Vegetables (except sugar beet)

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Alternaria leaf spot <i>Alternaria</i> spp.	3 to 5	15	7
	Cercospora leaf spot <i>Cercospora</i> spp.			
	Powdery mildew <i>Erysiphe</i> spp., <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp.			
<ul style="list-style-type: none"> <li>• <b>Spray Interval</b> - Apply before onset of disease and on a minimum interval of 7 days.</li> </ul>				

### Use Restrictions

- **DO NOT** apply more than 5 fl ozs (0.13 lb mefentrifluconazole) per acre per application.
- **DO NOT** make more than 3 applications at 5 fl ozs per acre per year.
- **DO NOT** apply more than 15 fl ozs (0.39 lb mefentrifluconazole) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products.

**Detailed Crop (Subgroup 1B) List** - beet, garden; burdock, edible; carrot; celeriac; chervil, turnip-rooted; chicory; ginseng; horseradish; parsley, turnip-rooted; parsnip; radish; radish, oriental (daikon); rutabaga; salsify; salsify, black; salsify, Spanish; skirret; turnip.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Cevya is a registered trademark of BASF.*

© 2021 BASF Corporation  
All rights reserved.

007969-00407.20210114.NVA 2020-04-549-0243  
Based on: NVA 2020-04-549-0145

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



We create chemistry