

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Hornet[®] WDG

EPA Reg. No. 62719-315

Tank Mixtures of Hornet[®] WDG herbicide and Callisto* herbicide plus Atrazine for Broad-Spectrum Weed Control in Field Corn

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Hornet[®] WDG broadleaf blend herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as described in this supplemental labeling, use of Hornet WDG is subject to all precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Hornet WDG may be tank mixed with Callisto* herbicide (mesotrione, 40% (w/w), 4 lb/gallon liquid) for broad-spectrum postemergence weed control in field corn. Apply a labeled rate of Hornet WDG with 0.5 to 3.0 fl oz of Callisto and 0.25 lb a.i./acre of atrazine. The combination of Hornet WDG and 0.25 lb a.i./acre of atrazine plus the minimum indicated rate of Callisto will control the weeds listed in the table below in addition to those weeds listed in the Postemergence Treatments section of the label for Hornet WDG.

Additional Weeds Controlled	Apply a Labeled Rate of Hornet WDG plus:	
	Callisto (fl oz/acre)	Atrazine (a.i./acre)
lambsquarters (1)	0.50	0.25
morningglory (annual)	0.75	0.25
nightshade	0.75	0.25
pigweed	0.75	0.25
waterhemp	0.75	0.25

1. Without atrazine, lambsquarters may be controlled with a labeled rate of Hornet WDG plus 1 fl oz/acre of Callisto.

Tank Mixing Restrictions: Do not exceed labeled rates of any tank mix product recommended in this supplemental labeling. Do not tank mix Hornet WDG with any product containing a label prohibition against tank mixing with products containing the active ingredient, flumetsulam. Refer to the General Information and Mixing sections, respectively, of the label for Hornet WDG for tank mixing precautions and tank mixing instructions. Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.

Application Timing: Apply as a broadcast postemergence treatment to field corn up to 20" tall or V6 stage, whichever occurs first. **Note:** Tank mixtures that contain atrazine are limited to 12" tall corn. For best results, apply when weeds are 2-4 inches in height and actively growing.

Required Adjuvants: Crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray) and either urea ammonium nitrate (UAN) at 2.5% v/v (2.5 gal/100 gal of spray) or spray grade ammonium sulfate (AMS) at 8.5 lb/100 gal of spray.

Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels. Use of this tank mix is permitted in only those states where both Hornet WDG and Callisto are registered.

®Trademark of Dow AgroSciences LLC
*Callisto is a trademark of Syngenta Crop Protection
R099-003
EPA Accepted: 01/13/03
Replaces D06-099-005.