

This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

S-METOLACHLOR GROUP 15 HERBICIDE

STRELIUS™ II



Contains S-metolachlor, the active ingredient used in Dual II Magnum®.

For Weed Control in Beans, Peas, and Lentils; Corn; Cotton; Grasses Grown for Seed; Horseradish; Peanuts; Potatoes; Pumpkin; Rhubarb; Safflowers; Sweet, Grain or Forage Sorghum; Soybeans; Soybeans, Immature Seed; Sugar Beets; Sugarcane*; Sunflowers; and Tomatoes.

***Not for Use in California Sale, Use and Distribution of this Product in Nassau and Suffolk Counties in the State of New York is Prohibited.**

ACTIVE INGREDIENT:	(% by weight)
S-metolachlor (CAS No. 87392-12-9)	82.4%
OTHER INGREDIENTS:	17.6%
TOTAL:	100.0%

Contains 7.64 lbs. of active ingredient per gallon. Formulated as an emulsifiable concentrate (EC).
EPA Reg. No.: 91234-50

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)
See below for additional Precautionary Statements.

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by a poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 - 20 minutes. ▪ Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> ▪ Move person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. ▪ Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

**For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night Within USA and Canada:
1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

Strelia™ II is not manufactured, or distributed by Syngenta Crop Protection, LLC, seller of Dual II Magnum®.



Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, or Viton \geq 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

Mixers and loaders supporting aerial applications are required to use closed systems. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)]. When using the closed system, the mixers' and loaders' PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Groundwater Advisory:

S-metolachlor is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory:

S-metolachlor has the potential to contaminate surface water through ground spray drift. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of S-metolachlor from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Non-Target Organism Advisory:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL AND CHEMICAL HAZARDS:

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

Mixing/Loading Instruction:

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsate.

Check-valves or anti-siphoning devices must be used on all mixing and/or irrigation equipment.

This product may not be mixed or loaded within 50 ft. of perennial or intermittent streams and rivers, natural or impounded lakes, and reservoirs. This product may not be mixed/loaded or used within 50 ft. of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Strelis II must be used only in accordance with directions on this label or in separately published EPA accepted supplemental labeling directions for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water are:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, or Viton \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

PRODUCT INFORMATION

Observe all precautions and limitations on the labels of each product used in tank mixtures. Tank mixtures are permitted only in those states where the tank mix partner is registered. Refer to and follow the label for each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

StreliaS II is a selective herbicide for use as a preplant surface-applied, preplant incorporated, or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in beans, peas, and lentils; corn (all types); cotton; grasses grown for seed; peanuts; potatoes; safflowers; sweet, grain, or forage sorghum; soybeans; soybean, immature seed; sugar beets; sugarcane; sunflowers; and tomatoes.

Restrictions: Do not use in nurseries, turf, or landscape plantings.

Do not apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas.

To prevent off-site movement due to runoff or wind erosion:

- A. Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
- B. Do not apply to impervious substrates, such as paved or highly compacted surfaces.
- C. Do not use tailwater from the first flood or furrow irrigation of treated fields to treat non-target crops, unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

Where directions specify a **StreliaS II** tank mixture with AAtrex[®] formulations, other brands of atrazine may be used. Follow the rates, recommendations, and limitations on the AAtrex or respective atrazine product label, if other brands of atrazine are used.

Restrictions: Certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

If **StreliaS II** is incorporated, any supplemental tillage before planting must not exceed the depth of incorporation.

Dry weather following preemergence application of **StreliaS II** or a tank mixture may reduce effectiveness. Cultivate if weeds develop.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control.

*Precaution: Injury may occur following the use of **StreliaS II** under abnormally high soil moisture conditions during early development of the crop.*

WEED RESISTANCE MANAGEMENT

For resistance management, **StreliaS II** is a Group 15 herbicide. Any weed population may contain or develop plants naturally resistant to **StreliaS II** and other Group 15 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **StreliaS II** or other Group 15 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available use the less resistance-prone partner at a rate that will control the target weeds equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation and that considers tillage or other mechanical control methods, cultural, biological and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds.
 - A spreading patch of non-controlled plants of a particular weed species.
 - Surviving plants mixed with controlled individuals of the same species.

If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance contact Atticus at 984-465-4800.

SOIL TEXTURES AND HERBICIDE RATES

Where rates are based on *coarse-, medium-, or fine-textured soils*, it is understood that soil textural classes are generally categorized as follows:

Coarse	Medium	Fine	
Sand	Loam	Sandy clay loam	Sandy clay
Loamy sand	Silt loam	Silty clay loam	Silty clay
Sandy loam	Silt	Clay loam	Clay

Within rate ranges in the rate tables and elsewhere on this label, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter.

Restrictions: **StreliaS II** may be applied preemergence alone, or in combination with tank mix partners specified on this label, following preplant incorporated herbicides when used according to their label directions, provided that such use is not prohibited on the respective labels.



Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. Do not use a sprayer or applicator contaminated with any other materials, or crop damage or clogging of the application device may result.

STRELIUS II APPLIED ALONE

Weeds Controlled:

Strelius II is taken up by the shoots and/or roots of emerging weeds. This uptake results in the inhibition of shoot and root tissue growth soon after weed germination. Because of this, **Strelius II** will not control emerged weeds and should be applied prior to weed emergence.

If **Strelius II** is incorporated, do not exceed a 2- to 3-inch depth. Any tillage after the **Strelius II** incorporation and before planting should not exceed 2 - 3 inches.

Dry weather following application of **Strelius II** may reduce weed control. Cultivate if weeds develop.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor or consistent control at a level below that generally considered acceptable for commercial weed control. Control of these weeds can be erratic, due partially to variable weather conditions. The following procedures may improve the control of weeds listed as partially controlled in **Table 1**:

- Thoroughly till soil to destroy germinating and emerged weeds.
- Plant crop into moist soil immediately after tillage. If **Strelius II** is to be used preemergence, apply at planting or immediately after planting.
- If available, sprinkler irrigate within 2 days after application. Apply 1/2 - 1 inch of water. Use lower water volume (1/2 inch) on *coarse-textured soils* and higher volume (1 inch) on *fine-textured soils*. Also, refer to the section on **CENTER PIVOT IRRIGATION APPLICATION** for this method of applying **Strelius II**.
- If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, a uniform, shallow cultivation is recommended as soon as weeds emerge.

Table 1: Weeds Controlled or Partially Controlled by Strelius II Applied Prior to Weed Emergence

Common Name	Scientific Name	Weed Type	Control (C) or Partial Control (PC)
Amaranth, Palmer	<i>Amaranthus palmeri</i>	Broadleaf	C
Amaranth, Powell	<i>Amaranthus powellii</i>	Broadleaf	C
Barnyardgrass	<i>Echinochloa crus-galli</i>	Grass	C
Beggarweed, Florida	<i>Desmodium tortuosum</i>	Broadleaf	PC
Carpetweed	<i>Mollugo verticillata</i>	Broadleaf	C
Crabgrass, large	<i>Digitaria ischaemum</i>	Grass	C
Crabgrass, smooth	<i>Digitaria sanguinalis</i>	Grass	C
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	Grass	C
Cupgrass, Prairie	<i>Eriochloa contracta</i>	Grass	C
Cupgrass, Southwestern	<i>Eriochloa acuminata</i>	Grass	C
Cupgrass, woolly	<i>Eriochloa villosa</i>	Grass	PC ¹
Eclipta	<i>Eclipta prostrata</i>	Broadleaf	PC
Foxtail, bristly	<i>Setaria verticillata</i>	Grass	C
Foxtail, giant	<i>Setaria faberi</i>	Grass	C
Foxtail, green	<i>Setaria viridis</i>	Grass	C
Foxtail, millet	<i>Setaria italica</i>	Grass	C
Foxtail, yellow	<i>Setaria pumila</i>	Grass	C
Galinsoga, hairy	<i>Galinsoga quadriradiata</i>	Broadleaf	C
Galinsoga, smallflower	<i>Galinsoga parviflora</i>	Broadleaf	C
Goosegrass	<i>Eleusine indica</i>	Grass	C
Johnsongrass (seedling)	<i>Sorghum halepense</i>	Grass	PC
Millet, wild-proso	<i>Panicum miliaceum</i>	Grass	PC ¹
Nightshade, Eastern black	<i>Solanum ptychanthum</i>	Broadleaf	C
Nightshade, hairy	<i>Solanum physalifolium</i>	Broadleaf	PC
Nutsedge, yellow	<i>Cyperus esculentus</i>	Sedge	C
Panicum, fall	<i>Panicum dichotomiflorum</i>	Grass	C
Panicum, Texas	<i>Panicum texanum</i>	Grass	PC
Pigweed, prostrate	<i>Amaranthus blitoides</i>	Broadleaf	C
Pigweed, redroot	<i>Amaranthus retroflexus</i>	Broadleaf	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	Broadleaf	C
Pigweed, tumble	<i>Amaranthus albus</i>	Broadleaf	C
Purslane, common	<i>Portulaca oleracea</i>	Broadleaf	PC
Pusley, Florida	<i>Richardia scabra</i>	Broadleaf	C
Rice, red	<i>Oryza punctata</i>	Grass	C
Ryegrass, Italian	<i>Lolium multiflorum</i>	Grass	C
Sandbur, field	<i>Cenchrus spinifex</i>	Grass	PC
Sandbur, Southern	<i>Cenchrus spinifex</i>	Grass	PC
Shattercane	<i>Sorghum bicolor</i>	Grass	PC
Signalgrass, broadleaf	<i>Urochloa platyphylla</i>	Grass	C
Sorghum (volunteer)	<i>Sorghum bicolor</i>	Grass	PC
Spiderwort, tropical	<i>Commelina benghalensis</i>	Broadleaf	C
Waterhemp, common	<i>Amaranthus rudis</i>	Broadleaf	C
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	Broadleaf	C
Witchgrass	<i>Panicum capillare</i>	Grass	C

¹ Refer to the **CORN** section of this label for additional directions.



REPLANT AND ROTATIONAL CROPS SECTION

Replanted Crop Directions:

This section covers replant crops that may be planted following a lost crop that has had an application of **Strelius II**.

If a crop treated with **Strelius II** is lost, any crop on this label, or on a supplemental **Strelius II** label, may be replanted immediately provided that the rate of **Strelius II** applied to the previous crop was not greater than the labeled rate for the crop to be replanted. If the first application was banded and the replant crop is planted in the center of the untreated bands, a second banded treatment may be applied at the rate for the use pattern for the replant crop, provided the application does not overlap the first application band.

Rotational Crop Directions:

Do not rotate to food or feed crops other than those listed below. For all crops not listed, wait at least 12 months following the last application of **Strelius II** before planting.

Barley, oats, rye, or wheat may be planted 4 1/2 months following treatment.

Alfalfa may be planted 4 months following application. Clover may be seeded 9 months following application.

If **Strelius II** is applied after June 1, rotating to crops other than corn (all types), grain, sorghum, or sugarcane the next spring, may result in crop injury.

Restrictions: To avoid injury to rotational alfalfa or clover, (1) Do not apply more than 1.9 lbs. active ingredient per acre (2.0 pts. of **Strelius II**) in the previous crop, and (2) Do not make lay-by or other postemergent applications of **Strelius II** in the previous crop.

Tobacco, buckwheat, rice, and sugarcane may be planted in the next spring following treatment.

Below in the rotational crop subsections **A through C** is a listing of rotational crop options that are made possible through *S*-metolachlor tolerances which were established by the EPA as crop groupings.

Precautions: 1) Rotating to crops within these crop groupings at less than 60 days may result in crop injury. 2) If the rate of **Strelius II** applied in the previous crops was greater than the rate listed here (Sections **A - C** below), these crops cannot be planted until the following spring.

1. If not more than 1.33 pts./A of **Strelius II** was applied to the field, the following crops (as well as those listed under subsections **B** or **C** below) may be planted 60 days after the last application. A second application of any *S*-metolachlor-containing product to the following crops is prohibited within 60 days of the original application:

- Crop Subgroup 1B - Root Vegetables:** garden beet, edible burdock, carrot, celeriac, turnip-rooted chervil, chicory, ginseng, horseradish, turnip-rooted parsley, parsnip, radish, oriental radish, rutabaga, salsify, black salsify, Spanish salsify, skirret, and turnip.
- Crop Subgroup 3 - Bulb Vegetables (if to be harvested green):** garlic, great-headed garlic, leek, green onion, Welsh onion, shallot.
- Winter squash (including pumpkins).

2. If not more than 1.67 pts./A of **Strelius II** was applied to the field, the following crops (as well as any listed under subsection **C** below) may be planted 60 days after the last application. A second application of an *S*-metolachlor-containing product to the following crops is prohibited within 60 days of the original applications:

- Crop Group 8 - Fruiting Vegetables, Except Cucurbits and Tabasco Peppers:** eggplant, groundcherry (*Physalis* spp.), pepino, peppers (bell, chili, cooking, pimento, and sweet), tomatillo, and tomato.

3. If not more than 2.0 pts./A of **Strelius II** was applied to the field, the following crops may be planted 60 days after the last application. A second application of an *S*-metolachlor-containing product to the following crops is prohibited within 60 days of the original application:

- Crop Subgroup 1C - Tuberos and Corm Vegetables:** arracacha; arrowroot; Chinese artichoke; Jerusalem artichoke; edible canna; bitter and sweet cassava; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; taniar; turmeric; yam bean; and yam, true.
- Crop Group 3 Bulb Vegetables (if to be harvested dry):** garlic, great-headed garlic, leek, dry bulb and green onion, Welsh onion, shallot.
- Crop Subgroup 4B - Leaf Petiole Vegetables:** cardoon, celery, Chinese celery, celtuce, Florence fennel, rhubarb, and Swiss chard.
- Crop Subgroup 5A - Head and Stem Brassica Vegetables:** broccoli; Chinese broccoli, Brussels sprouts, cabbage, Chinese (Napa) cabbage, Chinese mustard, cauliflower, cavalo broccolo, and kohlrabi.

APPLICATION PROCEDURES

Application Timing: **Strelius II** alone or in some tank mixtures with other labeled herbicides may be applied for weed control in certain crops at various times. Refer to the given crop section of the label to determine if application timings listed below are permitted. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixtures.

Preplant Surface-applied: For minimum-tillage or no-tillage systems only, **Strelius II** alone and some **Strelius II** tank mixtures may be applied up to 45 days before planting certain crops. Use only split applications for treatments made 30 - 45 days before planting, with 2/3 the recommended broadcast rate for the crop and soil texture applied initially and the remaining 1/3 at planting. Treatments less than 30 days before planting may be made either as a split or a single application. Refer to individual crop section on this label to determine if early preplant surface application is permitted. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (for example, Gramoxone® brands, Touchdown® brands, or Roundup® brands). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated: Apply **Strelius II** to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate **Strelius II** after bed formation, unless specified otherwise.

Preemergence: Apply **Strelius II** during planting (behind the planter) or after planting, but before weeds or crops emerge.

Postemergence: **Strelius II** will not control emerged weeds so it must be applied to a weed-free soil surface or in tank mixture with products that provide postemergence control of weeds present at the time of application. Refer to the individual crop section of this label if a postemergence application is permitted.

SPECIAL APPLICATION PROCEDURES

CA Only (Beans, Peas, and Lentils; Corn; Safflowers):

Preplant Incorporated: Broadcast **Strelius II** alone or with tank mix partners listed on this label to the soil and thoroughly incorporate with a disk or similar implement set to till 4 - 6 inches deep. For more thorough incorporation, till the soil in 2 different directions (cross-till). Crops may be planted on flat surface or on beds. Use caution when forming the beds that only soil from the **Strelius II** treated zone is used (i.e., untreated soil should not be brought to soil surface). If the application is made to preformed beds, incorporate **Strelius II** with a tillage implement set to till 2 - 4 inches deep. Care should be taken during tilling to keep the tilled (**Strelius II**-treated) soil on the beds.

Preemergence: Apply **Strelius II** after planting. Water with sprinkler or flood irrigation within 7 - 10 days.

Fall Application for Spring Weed Control

(Only in IA, MN, ND, SD, WI, and portions of NE and IL - See specific instructions in the BEANS, PEAS, AND LENTILS; CORN; and SOYBEANS sections of this label for timing of application and other information):

Do not apply to frozen ground. Use on *medium* and *fine* soils with greater than 2.5% organic matter that will be planted to corn or soybeans the next spring. Ground may be tilled before or after application. Do not exceed a 2 to 3-inch incorporation depth if tilled after treatment.

Restrictions: If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for the specific crop, or illegal residues may result.

Fall Application for Italian Ryegrass Control

(Corn, Cotton, Grain and Forage Sorghum, and Soybean Only - See specific instructions in the CORN, COTTON, GRAIN AND FORAGE SORGHUM, and SOYBEAN sections of this label for timing of application and other information):



Strelius II may be applied in the fall (September 1 - December 1) for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). A tillage operation may precede the application. Do not incorporate to a depth greater than 2 - 3 inches if tillage follows the application of **Strelius II**.

- Restrictions:**
1. Do not apply **Strelius II** to frozen ground.
 2. All crops on the **Strelius II** label may be planted the following spring after application.
 3. If a spring application is made, the combined total amount of **Strelius II** applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for the specific crop planted, or illegal residues may result. Refer to the crop sections on this label for specific directions.

Ground Application:

Apply **Strelius II** alone or in tank mixtures by ground equipment in a minimum of 10 gals. of spray mixture per acre, unless otherwise specified. Use sprayers that provide accurate and uniform application. For **Strelius II** tank mixtures with wettable powder or dry flowable formulations, screens and strainers should be no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after use. Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

- For information on applying in lower volumes of carrier, see **LOW CARRIER APPLICATION** section.
For application by air or through center pivot systems, see **Mandatory Spray Drift Management** and **Center Pivot Irrigation Application** sections.
For information on impregnating dry fertilizer, see **DRY BULK GRANULAR FERTILIZER** section.
For information on application using variable-rate technologies, see **Variable-rate Application** section.

SPRAY EQUIPMENT

LOW CARRIER APPLICATION

For Broadcast Ground Application Only:

Use sprayers, such as Ag-Chem RoGator®, Hagie, John Deere Hi-Cycle™, Melroe Spra-Coupe, Tyler Patriot™, or Willmar Air Ride®, that provide accurate and uniform application. **Only water may be used as a carrier.** Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens as fine as 100-mesh be used with some nozzles. Use a pump with capacity to: (1) maintain up to 35 - 40 psi at the nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension. Use a minimum of 5.0 gals. of spray mixture per acre. Maximum recommended sprayer speed is 15 mph. Rinse sprayer thoroughly with clean water immediately after each use. **Precautions:** Low pressure nozzles are recommended to reduce drift and increase application accuracy. Care should be taken when using automatic rate controlling devices to spray the material within the rated working pressure and flow ranges of the nozzles selected. Nozzle screens should be used when recommended by the manufacturer. All nozzles should be placed on 20-inch centers, except flooding types which should be placed on 40-inch centers. When Flat Fan-type nozzles are used, angles of 80° or 110° are recommended. Always read and follow the manufacturer's directions for optimum setup and performance of their nozzles or tips.

Aerial Application:

Apply **Strelius II** in water alone or in tank mixtures with AAtrex, Lorox®, or Sencor® in a minimum total volume of 2.0 gals./A by aircraft. **Strelius II** may also be applied by air in combination with Prowl® or Treflan®. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 15 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply **Strelius II** alone or **Strelius II** + AAtrex by aircraft at a minimum upwind distance of 400 ft. from sensitive plants, or apply **Strelius II** + Lorox or Sencor at a minimum upwind distance of 300 ft. from sensitive plants.

The applicator must be familiar with and take into account the information covered in the **Mandatory Spray Drift Management** and **Spray Drift Advisories** sections below.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select the nozzle and pressure that deliver medium or coarser droplets (ASABE S641).
- If the wind speed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the field. When the wind speed is between 11-15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select the nozzles and pressure that deliver medium or coarser droplets (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to select the nozzle and pressure that deliver medium or coarser droplet size (ASABE S572) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.



Controlling Droplet Size – Aircraft

• **Adjust Nozzles** - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

BOOMLESS GROUND APPLICATIONS:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

HANDHELD TECHNOLOGY APPLICATIONS:

Take precautions to minimize spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

CENTER PIVOT IRRIGATION APPLICATION

StreliaS II alone or in tank mixture with other herbicides on this label, which are registered for center pivot application, may be applied in irrigation water preemergence (after planting, but before weeds or crop emerge) at rates listed on this label. **StreliaS II** also may be applied postemergence to the crop and preemergence to weeds in crops where postemergence applications are allowed on this label. Follow all restrictions (height, timing, rate, etc.) to avoid illegal residues. Apply this product only through a center pivot irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Operating Instructions:

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Prepare a mixture with a minimum of 1-part water to 1-part herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
9. Meter into irrigation water during entire period of water application.
10. Apply in 1/2 - 1 inch of water. Use the lower water volume (1/2 inch) on *coarse-textured soils* and the higher volume (1 inch) on *fine-textured soils*. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

Precaution for Center Pivot Applications: Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.

DRY BULK GRANULAR FERTILIZERS

Many dry bulk granular fertilizers may be impregnated or coated with **StreliaS II** alone or selected **StreliaS II** tank mixtures which are registered for preplant incorporated or preplant surface applications which are used to control weeds in crops on the **StreliaS II** label and are not prohibited from use on dry bulk granular fertilizers.

When applying **StreliaS II** or **StreliaS II** mixtures with dry bulk granular fertilizers, follow all directions for use and precautions on the respective product labels, regarding target crops, rates per acre, soil texture, application methods (including timing of application), and rotational crops.

All individual state regulations relating to dry bulk granular fertilizers blending, registration, labeling, and application are the responsibility of the individual and/or company selling the herbicide/fertilizer mixture.

Prepare the herbicide/fertilizer mixtures by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray **StreliaS II** and **StreliaS II** mixtures onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer only and to avoid spraying the walls of the blender.

If the herbicide/fertilizer mixture is too wet, add a highly absorptive material, such as Agsorb® or Celatom MP-79®, or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture. Add absorptive materials only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application results will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer material being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight.

Calculate amounts of **StreliaS II**, AAtrex, AAtrex + Princep®, Balance® Pro, Princep, Sencor, or Sonalan® by the following formula:

$$\frac{2,000}{\text{lbs. of fertilizer per acre}} \times \text{pts./A of liquid or flowable product} = \text{pts. of liquid or flowable product per ton of fertilizer}$$
$$\frac{2,000}{\text{lbs. of fertilizer per acre}} \times \text{pts./A of dry product} = \text{pts. of dry product per ton of fertilizer}$$



Pneumatic (Compressed Air) Application (Strelius II Alone):

High humidity, high urea concentrations, low fertilizer use rates, and dusty fertilizer may cause fertilizer mixture to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, premix **Strelius II** with Exxon Aromatic 200 at a rate of 1.0 - 4.0 pts./gals. of **Strelius II**. Aromatic 200 is a noncombustible/nonflammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents should not be used when using Aromatic 200.

Restriction:

1. Mixtures of **Strelius II** and Aromatic 200 must be used on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water or liquid fertilizer solutions for spraying applications.

Precautions:

1. To avoid potential for explosion, do not impregnate **Strelius II** or **Strelius II** mixtures on ammonium nitrate, potassium nitrate, or sodium nitrate, either alone or in blends with other fertilizers.
2. To avoid potential for explosion, do not use **Strelius II** or **Strelius II** mixtures on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.
3. Drying agents are not recommended for use with On-The-Go impregnation equipment.
4. When impregnating **Strelius II** in a blender before application, a drier mixture can be attained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or drying agents of 6/30 particle size are recommended.

Application:

Apply 200 - 700 lbs. of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Nonuniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control. On *fine- or medium-textured soils* in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, make applications approximately 30 days before planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On *coarse-textured soils*, make applications approximately 14 days prior to planting.

Precaution: To avoid crop injury, do not use the herbicide/fertilizer mixture on crops where bedding occurs.

MIXING INSTRUCTIONS

Strelius II Alone:

Mix **Strelius II** with water or fluid fertilizer and apply as a spray. Fill the spray tank 1/2 - 3/4 full with water or fluid fertilizer, add the proper amount of **Strelius II**, then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

Tank Mixtures:

Fill the spray tank 1/4 full with water, and start agitation; add 2,4-D, AAtrex, Balance Pro, Banvel[®], Basagran[®], Butyrac[®], Canopy[®], Caparol[®] 4L, Command[®], Cotoran[®], Eptam[®], Liberty[®] Herbicide, Lorox, Marksman[®], MSMA, Princep, Prowl, Pursuit[®], AAtrex + Princep, Scepter[®], Sencor, Sonalan, or Treflan, and allow it to become dispersed; then add **Strelius II**; then add Gramoxone brands, Landmaster[®] BW, or Touchdown or Roundup (glyphosate products) if these products are being used; and finally the rest of the water. For tank mixtures with AAtrex, Balance, Banvel, Canopy, Caparol 4L, Command, Cotoran[®], Eptam, Lorox, Marksman, Princep, Prowl[®], Pursuit, AAtrex + Princep, Scepter, Sencor, Sonalan, or Treflan, fluid fertilizers may replace all or part of the water as carrier, except in the AAtrex postemergence and the Banvel postemergence tank mixes. For tank mixtures with AAtrex, see additional mixing instructions on the AAtrex label. For each mixture, check compatibility with fluid fertilizer, as described below, before mixing in spray tank. For all tank mixtures, agitate during mixing and application to maintain a uniform suspension.

* See **Special Mixing Instructions** for tank mixtures with Cotoran and with AAtrex or Princep + Prowl under the appropriate tank mixture section.

For directions on how to conduct a compatibility test, see the **COMPATIBILITY TEST** section.

COMPATIBILITY TEST

A jar test is recommended before tank mixing to ensure compatibility of **Strelius II** with other pesticides. The following test assumes a spray volume of 25 gals./A. For other spray volumes, make appropriate changes in the ingredients.

Restrictions: Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray. Because liquid fertilizers vary, even within the same analysis, **always check compatibility with pesticide(s) before use.** Incompatibility of tank mixtures is more common with suspensions of fertilizer and pesticides.

Test Procedure:

1. Add 1.0 pt. of carrier (fertilizer or water) to each of 2 one qt. jars with tight lids. **Restrictions:** Use the same source of water that will be used for the tank mix and conduct the test at the temperature the tank mix will be applied.
2. To one of the jars, add 1/4 tsp. or 1.2 milliliters of a compatibility agent approved for this use, such as Compex[®] or Unite[®] (1/4 tsp. is equivalent to 2.0 pts./100 gals. spray). Shake or stir gently to mix.
3. To both jars, add the appropriate amount of pesticide(s) in their relative proportions based on specified label rates. If more than one pesticide is used, add them separately with dry pesticides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix.
4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixtures stand 15 - 30 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (a) Slurry the dry pesticide(s) in water before addition, or (b) add 1/2 the compatibility agent to the fertilizer or water and the other 1/2 to the emulsifiable concentrate or flowable pesticide before addition to the mixture. If incompatibility is still observed, do not use the mixture.
5. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the **STORAGE AND DISPOSAL** section in this label.

CROP USE DIRECTIONS

CORN (ALL TYPES)

STRELIUS II ALONE

Apply **Strelius II**, either preplant surface, preplant incorporated, preemergence, postemergence, or lay-by, using the appropriate rate specified below.

Preplant Surface-Applied:

Refer to instructions for use of **Strelius II** alone under **APPLICATION PROCEDURES**.

Fall Application for Spring Weed Control:

1. Apply after September 30 in ND, SD, MN, WI and north of Route 30 in IA.
2. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
3. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 - 2.0 pts./A on *medium-textured* and 2.0 pts./A on *fine-textured soils*. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2 - 3 inches. Minimize furrow and ridge formation in the tillage operations.

Restrictions: If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for corn, or illegal residues may result.



Fall Application for Italian Ryegrass Control:

StreliaS II may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply **StreliaS II** at 1.33 - 1.67 pts./A in the fall (September 1 - December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower **StreliaS II** rate for *coarse-textured soils* and the higher rate for *fine-textured soils*. A tillage operation may precede the application. Do not incorporate to a depth greater than 2 - 3 inches if tillage follows the application of **StreliaS II**. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone brands can be tank mixed with **StreliaS II** to control emerged ryegrass. Refer to the Gramoxone brands label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with **StreliaS II** for control or improved control of other weeds present at the time of application.

Restrictions:

1. Do not apply **StreliaS II** to frozen ground.
2. If a spring application is made, the combined total amount of **StreliaS II** applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for corn (3.9 pts./A, depending on soil texture) or illegal residues may result.

Fall Application for Control or Suppression of Yellow Nutsedge (ID, OR, and WA only):

For preemergent control or suppression of yellow nutsedge the following spring, apply 1.33 pts./A of **StreliaS II** in the fall after the harvest of the previous crop but before freeze-up. Fall applications of **StreliaS II** can be surface-applied or incorporated.

Restrictions:

1. Make no more than one fall application per crop.
2. Apply not more than 1.33 pts./A in a single fall preplant application.
3. Do not apply to frozen ground.
4. If a spring application is made, the combined total amount of **StreliaS II** applied in the fall plus the spring must not exceed the maximum season S-metolachlor rate for corn (3.9 pts./A, depending on soil texture), or illegal residues may result.

EARLY PREPLANT APPLICATIONS

Use on *medium-* and *fine-textured soils* with minimum-tillage or no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply 2/3 the labeled rate of **StreliaS II** (1.67 pts./A on *medium soils* and 2.0 pts./A on *fine soils*) as a split treatment 30 - 45 days before planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Apply 1.33 pts./A on *coarse soils* not more than 2 weeks before planting.

Restrictions: If a spring application is made, the total rate of the fall plus spring application must not exceed the maximum total rate for corn, or illegal residues may result.

On *medium-* and *fine-textured soils* with minimum- or no-tillage systems in CT, DE, MA, MD, ME, MI, NH, NY, OH, PA, RI, VA, VT, and WV, preplant surface applications may be applied following the directions for use above. If the amount of rainfall results in unsatisfactory length of weed control following the earlier treatment, a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide may be used, i.e., AAtrex, Beacon®, Bicep Magnum®, Bicep II Magnum®, Exceed®, Accent®, Banvel, Basagran, bromoxynil (Buctril®), or 2,4-D. If the postemergence treatment includes the herbicide used preplant surface-applied, do not exceed the total labeled rate for corn on a given soil texture. Observe all directions for use, precautions, and limitations on the label of the postemergent herbicide.

PREPLANT INCORPORATED OR PREEMERGENCE

Follow instructions for use of **StreliaS II** alone under **APPLICATION PROCEDURES**. On *coarse soils*, apply 1.0 - 1.33 pts./A of **StreliaS II** if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts./A of **StreliaS II**. On *fine soils*, apply 1.33 - 1.67 pts./A of **StreliaS II** if organic matter content is less than 3%, or 1.67 - 2.0 pts./A if organic matter content is 3% or greater.

POSTEMERGENCE OR LAY-BY

To extend the duration of weed control in corn, a maximum rate of 2.0 pts./A of **StreliaS II** may be applied after corn emergence until the corn plants reach 40 inches in height, following any preplant surface-applied, preplant incorporated, or preemergence herbicide application, including **StreliaS II**. For best results, applications should be made to soil free of merged weeds and directed toward the base of corn plants in excess of 5 inches tall. The total **StreliaS II** rate applied on corn during any one crop year should not exceed 3.9 pts./A, depending on soil texture.

Restrictions for All Applications to Corn: To avoid possible illegal residues:

1. Do not graze or feed forage from treated areas for 30 days following application.
2. Do not harvest sweet corn ears from treated areas for 30 days following application.

PROBLEM WEED CONTROL DIRECTIONS

Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta - Partial Control:

For more consistent partial control of shattercane, wild proso millet, woolly cupgrass, or eclipta, apply 1.0 - 1.33 pts./A of **StreliaS II** preplant incorporated followed by 1.0 - 1.33 pts./A of **StreliaS II** preemergence. Make the preemergence application during or after planting, but before weeds and corn emerge. Apply the 1.33 pts./A rate of **StreliaS II** when a heavy infestation of shattercane, wild proso millet, woolly cupgrass, or eclipta is expected. A shallow cultivation may be needed to control any late emerging weeds.

Woolly Cupgrass and Wild Proso Millet Control Program:

For control of these species, use the following 3-step program:

1. Apply **StreliaS II** early preplant, preplant incorporated, or preemergence at 1.67 pts./A on *medium soils* and 2.0 pts./A on *fine-textured soils*, up to the maximum label rate. Lightly incorporate with a rotary hoe if rainfall does not occur within 5 - 7 days.
2. Apply a postemergence tank mix of Beacon at 0.38 oz./A or Exceed at 1 packet per 4 acres plus Accent SP at 0.33 oz./A plus 1.0 qt. of crop oil concentrate plus 1.0 gal./A of 28% nitrogen, or the equivalent amount of ammonium sulfate, when grasses are 2 - 3 inches tall and the corn is at least 4 inches tall.
3. Cultivate 14 - 21 days after the postemergence application.

Restrictions:

1. Do not apply more than the labeled application rate for a given soil texture per year, either as a single or split treatment, or illegal residues may result.
2. In corn, use up to 2.5 pts./A of **StreliaS II** preplant surface, preplant incorporated, or preemergence treatment on soils having an organic matter content between 6% and 20%.
3. In the event of escape of annual weeds following a preplant surface, preplant incorporated, or preemergence treatment of **StreliaS II**, follow with a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide, i.e., AAtrex, Beacon, Bicep II Magnum, Accent, Banvel, Basagran, Buctril, or 2,4-D. If the postemergence treatment includes the herbicide used in the earlier treatment, do not exceed the total labeled rate for corn on a given soil texture.
4. Buctril may be applied postemergence alone or in tank mix combination with AAtrex. Do not exceed 1.2 lbs. a.i./A of AAtrex in tank mix combination with Buctril postemergence. Refer to the AAtrex and Buctril labels for specific rates and precautions.
5. Do not use **StreliaS II** on peat or muck soils.



STRELIUS II COMBINATIONS

StreliaS II in any tank mixture for corn may be applied in water or fluid fertilizer before corn emerges. Use only water as a carrier when **StreliaS II** is applied after corn emergence.

Restrictions: For all applications to corn:

1. Do not graze or feed forage from treated areas for 30 days following application.
2. Do not harvest sweet corn ears from treated areas for 30 days following application or possible illegal residues may result.

IMPORTANT: FOR TANK MIXTURES WITH AATREX (OR OTHER BRANDS OF ATRAZINE) - IF APPLYING StreliaS II IN TANK MIXTURE WITH AATREX, ALL THE RESTRICTIONS AND RATE LIMITATIONS ON THE AATREX LABEL MUST BE FOLLOWED IF MORE RESTRICTIVE/PROTECTIVE THAN THOSE ON THIS LABEL. IN ADDITION, IF AATREX IS/MUST BE APPLIED AT RATES LOWER THAN THOSE LISTED ON THIS LABEL, BROADLEAF WEED CONTROL MAY BE AFFECTED. REFER TO THE AATREX LABEL FOR WEEDS CONTROLLED AT THE REDUCED RATES.

Table 2: StreliaS II Tank Mixtures for Corn - Additional Weeds Controlled and Special Instructions

	StreliaS II + AAtrex and/or Princep (Preplant Surface, PPI, PRE)	StreliaS II + AAtrex (Post)	StreliaS II + Banvel (Field Corn)	StreliaS II + AAtrex + Lorax	StreliaS II + AAtrex or Princep + Prowl	StreliaS II + Marksman	StreliaS II + Balance Pro
Special Mixing Instructions					1		
Comments	2, 3, 4, 5, 7, 8	2, 3, 4, 5		2, 3, 4, 5, 6	2, 3, 4, 5	7	2, 3, 7
Browntop panicum	X			X	X		X
Cocklebur	X	0	0	X	X		0 - X
Common purslane	X			X	X	X	X
Hairy nightshade	X			X	X		X
Jimsonweed		X	0			X	X
Kochia		X				X	X
Lambsquarters	X	X	X	X	X	X	X
Morningglory	X	0	0	X	X		X
Mustard		X				X	X
Pigweed				X	X	X	X
Prickly sida		X					
Ragweed	X	X	X	X	X	X	X
Smartweed	X	X	X	X	X	X	X
Velvetleaf	X	X	0	X	X	0 - X	0 - X

X = control; 0 = partial control; 0 - X = partial to full control depending on ratio of products used or on weed population.

Comments:

1. Special Mixing Instructions for **StreliaS II** + AAtrex or Princep and Prowl
 - a. Fill the spray tank 1/4 full with water or fluid fertilizer and start agitation.
 - b. To aid compatibility, add a compatibility agent, such as Unite or X-77®, at 4.0 pts./100 gals. of spray mixture.
 - c. Then add the AAtrex or Princep and allow it to become dispersed.
 - d. Then add **StreliaS II** and Prowl 4E.
 - e. Add the rest of the water.
2. Although a single formulation for AAtrex or Princep is listed in the rate tables, other formulations may be substituted, using the following formula:
 - a. 1.0 lb. of AAtrex® Nine-0® or Princep® Caliber 90® = 1.8 pts. of AAtrex 4L or Princep 4L
3. Although direction specify AAtrex formulations in tank mixture with **StreliaS II**, other brands of atrazine may be used. Follow the rates, recommendations, and limitations on the atrazine label.
4. See additional mixing instructions on the AAtrex label.
5. *Precaution: Do not exceed a total of 2.5 lbs. a.i. of atrazine per acre per year. However, certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.*
6. Other formulations of Lorax can be used: 1.0 lb. of Lorax DF = 1.0 pt. of Lorax L.
7. In minimum-tillage and no-tillage systems, mix with Gramoxone brands for control of most emerged annual weeds and suppression of perennial weeds; or with Landmaster BW for suppression of emerged field bindweed and control or suppression of annual weeds; or with Touchdown brands or Roundup brands for control of most emerged annual and perennial weeds.
8. Refer to the **CORN - STRELIUS II COMBINATIONS - TANK MIXTURE WITH AATREX; OR AATREX + 2,4-D; OR AATREX + 2,4-D + BANVEL FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS** sections for specific directions for 2,4-D or Banvel burndown combinations in minimum-tillage or no-tillage systems.

StreliaS II in any tank mixture for corn may be applied in water or fluid fertilizer, except as noted.

Restrictions:

1. For all applications to corn, do not graze or feed forage from treated areas for 30 days following application and do not harvest sweet corn ears from treated areas for 30 days following application, or possible illegal residues may result.
 2. When applying **StreliaS II** in tank mixture with AAtrex, do not exceed a total of 2.5 lbs. a.i. of atrazine per acre per year.
- Refer to **CORN (ALL TYPES) - STRELIUS II ALONE** for directions on sequential postemergence treatments if escape weeds develop.

TANK MIXTURE WITH AATREX OR PRINCEP, OR AATREX + PRINCEP - PREPLANT SURFACE, PREPLANT INCORPORATED, OR PREEMERGENCE

In addition to the weeds controlled by **StreliaS II** alone, **StreliaS II** + AAtrex or Princep, or **StreliaS II** + AAtrex + Princep, applied preplant surface, preplant incorporated, or preemergence, also controls the following weeds: browntop panicum, cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

Apply **StreliaS II** + AAtrex or Princep, or **StreliaS II** + AAtrex + Princep either preplant surface, preplant incorporated, or preemergence.



Preplant Surface-applied:

Follow instructions for use of **Strelius II** alone under **APPLICATION PROCEDURES** and under application instructions for **STRELIUS II ALONE** on corn. Apply **Strelius II** + AAtrex or Princep, or **Strelius II** + AAtrex + Princep on *medium soils* (1.67 pts./A of **Strelius II** + 3.2 - 4.0 pts./A of AAtrex 4L or Princep 4L, or AAtrex 4L + Princep 4L combined) and on *fine soils* (1.67 - 2.0 pts./A of **Strelius II** + 4.0 pts./A of AAtrex 4L or 4.0 - 5.0 pts./A of Princep 4L, or AAtrex 4L + Princep 4L combined) in minimum-tillage and no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply the tank mixtures as a split or single treatment in those states and as indicated in the **STRELIUS II ALONE - Preplant Surface-applied** section of the label for corn. On *coarse soils*, apply 1.33 pts./A of **Strelius II** and 3.2 pts./A of AAtrex 4L or Princep 4L, or AAtrex 4L + Princep 4L combined.

Preplant Incorporated or Preemergence:

Follow instructions for use of **STRELIUS II** alone under **APPLICATION PROCEDURES**. Apply **Strelius II** + AAtrex or Princep, or **Strelius II** + AAtrex + Princep, using the appropriate rates from **Table 3**.

Restrictions: Do not apply more than the labeled rate for a given soil texture per year, either as a split or single treatment, or illegal residues may result.

Shattercane and Wild Proso Millet - Partial Control

For more consistent partial control of shattercane or wild proso millet where **Strelius II** is applied in tank mixture or sequentially with other registered corn herbicides, the following applications may be made:

1. Apply 1.0 - 1.33 pts./A of **Strelius II** + 2.0 lbs. a.i./A of AAtrex or Princep preplant incorporated, followed by 1.0 - 1.33 pts./A of **Strelius II** preemergence. Make the preemergence application during or after planting, but before weeds and corn emerge.
2. Apply **Strelius II** at 1.33 pts./A alone or in tank mix combination with up to 2.0 lbs. a.i./A of AAtrex, or Princep, preplant incorporated. Do not exceed the total rate of triazine herbicide labeled in combination with **Strelius II** for corn grown on a given soil texture. Follow with a post-directed application of Evik® 80 WDG. Refer to the Evik 80 WDG label for specific directions for the post-directed application.
3. Apply Eradicane® (or equivalent EPTC or butylate formulations) at labeled rates preplant incorporated, followed by a preemergence application of **Strelius II** at 1.0 - 1.33 pts./A. Do not use Eradicane on soils where rapid degradation has been shown to occur. Make the preemergence application during or after planting, but before weeds and corn emerge.

Precautions: When following the application regimes in numbers 1 - 3 above, a shallow cultivation may be needed after the preemergence or postemergence application to help control any late emerging shattercane or wild proso millet plants.

Restrictions: Do not exceed a total of 1.9 lbs. a.i./A (2.0 pts. of **Strelius II**) in the preplant incorporated plus preemergence application on soils with less than 6% organic matter, or crop injury may result.

Table 3: Strelius II + AAtrex or Princep, or Strelius II + AAtrex + Princep, Preplant Incorporated or Preemergence - Corn (All Types)

Soil Texture	Broadcast Rates Per Acre					
	Less than 3% Organic Matter			3% Organic Matter or Greater		
	Strelius II + AAtrex Nine-0* or Princep Caliber 90*	OR	Strelius II + AAtrex Nine-0** + Princep Caliber 90**	Strelius II + AAtrex Nine-0* or Princep Caliber 90*	OR	Strelius II + AAtrex Nine-0** + Princep Caliber 90**
COARSE	0.8 - 1.0 pt. + 1.1 - 2.2 lbs.		0.8 - 1.0 pt. + 0.6 - 1.1 lbs. + 0.6 - 1.1 lbs.	1.0 pt. + 1.3 - 2.2 lbs.		1.0 pt. + 0.7 - 1.1 lbs. + 0.7 - 1.1 lbs.
MEDIUM	1.0 - 1.33 pts. + 1.3 - 2.2 lbs.		1.0 - 1.33 pts. + 0.7 - 1.1 lbs. + 0.7 - 1.1 lbs.	1.33 pts. + 1.8 - 2.2 lbs.		1.33 pts. + 0.9 - 1.1 lbs. + 0.9 - 1.1 lbs.
FINE	1.33 pts. + 1.8 - 2.2 lbs.		1.33 pts. + 0.9 - 1.1 lbs. + 0.9 - 1.1 lbs.	1.33 - 1.67 pts. + 1.8 - 2.2 lbs.***		1.33 - 1.67 pts. + 0.9 - 1.1 lbs.*** + 0.9 - 1.1 lbs.***
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE					

* Use Princep in preference to AAtrex when heavy infestations of crabgrass or fall panicum are expected. On soils having between 6% and 20% organic matter, **Strelius II** may be used up to 2.33 pts./A in tank mix combination with 2.2 lbs./A of AAtrex Nine-0, or equivalent rates of AAtrex 4L. Refer to the AAtrex label for weeds controlled at this reduced rate.

** When using the tank mixture of **Strelius II** + AAtrex Nine-0 + Princep Caliber 90, use equal rates of each as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of AAtrex + Princep instead of the 1:1 ratio given in **Table 3**. (Example: Total AAtrex Nine-0 + Princep Caliber 90 = 1.2 lbs./A, use 0.4 lb. of AAtrex + 0.8 lb. of Princep, respectively.) Refer to **Comment No. 2** following **Table 2** for AAtrex 4L and Princep 4L conversions.

*** For cocklebur, yellow nutsedge, and velvetleaf control on *fine-textured soils* above 3% organic matter, apply 2.25 lbs./A of AAtrex Nine-0, or equivalent rates of AAtrex 4L, or the same total amount of AAtrex + Princep with 1.33 - 1.67 pts./A of **Strelius II**.



TANK MIXTURE WITH AATREX - POSTEMERGENCE

Weeds Controlled		
Barnyardgrass (watergrass)	Green foxtail	Prickly sida
Crabgrass	Jimsonweed	Purslane
Crowfootgrass	Kochia	Ragweed
Fall panicum	Lambsquarters	Smartweed
Giant foxtail	Mustard	Velvetleaf
	Pigweed	Yellow foxtail
Weeds Partially Controlled		
Cocklebur	Morningglory	Yellow nutsedge

Apply 1.0 pt./A of **StreliaS II** + 1.3 lbs./A of AAtrex Nine-0* on *coarse soils*, 1.33 pts./A of **StreliaS II** + 1.8 lbs./A of AAtrex Nine-0 on *medium soils*, or 1.33 - 1.67 pts./A of **StreliaS II** + 1.8 - 2.2 lbs./A** of AAtrex Nine-0 on *fine soils*. Apply this tank mixture before grass and broadleaf weeds pass the 2-leaf stage and before corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally result in unsatisfactory control.

Lay-by:

Apply to corn plants not more than 12 inches tall. Applications to corn in excess of 5 inches should be directed to the base of the corn plants; whereas, applications to corn plants less than 5 inches tall may be made over the top. Occasionally, some corn leaf burn may result, but this should not affect later growth or yield. Do not apply this postemergence tank mixture in fluid fertilizer, or severe crop injury may occur.

*When using AAtrex 4L, use equivalent rates. One lb. of AAtrex Nine-0 = 1.8 pts. of AAtrex 4L.

For better control of cocklebur, morningglory, velvetleaf, and yellow Nutsedge on *fine-textured soils* above 3% organic matter, apply 2.2 lbs./A of AAtrex Nine-0, or equivalent rate of AAtrex 4L with 1.33 - 1.67 pts./A of **StreliaS II. Tank mixtures of **StreliaS II** + AAtrex may be applied following use of any registered preplant surface-applied, preplant incorporated, or preemergence corn herbicide, including **StreliaS II** + AAtrex.

Restrictions: The total **StreliaS II** rate should not exceed 3.9 pts., nor the AAtrex rate more than 2.5 lbs. a.i./A during any one crop year, or illegal residues may result. Refer to the AAtrex label for geographic, soil-texture, and rotational restrictions.

TANK MIXTURE WITH BANVEL

Preemergence:

Use this tank mixture only on field corn which is flat-planted (no furrows) in CO, IA, IL, IN, KS, MN, NE, OH, SD, and WI.

In addition to the weeds controlled by **StreliaS II** alone, **StreliaS II** + Banvel, applied preemergence, also controls lambsquarters, ragweed, smartweed, cocklebur*, jimsonweed*, morningglory*, and velvetleaf*.

*Partially controlled.

Apply **StreliaS II** + Banvel preemergence. Broadcast 1.0 pt./A of Banvel with 1.33 pts./A of **StreliaS II** on *medium soils*, or with 1.33 - 1.67 pts./A of **StreliaS II** on *fine soils*. Do not apply on *coarse soils* or on soils with less than 2.5% organic matter. Apply this tank mixture to the soil surface at planting or after planting, but before corn emerges. Plant corn at least 1.5 inches deep and apply behind planting equipment, avoiding incorporation by the planter wheel or other seed covering device. Do not incorporate before corn emergence. If it is necessary to rotary hoe to break the soil crust, do not disturb the soil more than 1/2 inch deep.

Restriction:

1. Do not apply with aircraft.

Precaution:

1. Avoid drift to sensitive nontarget plants, such as soybeans, during application, or injury may occur.

Postemergence for Control of Pigweed (Mid-atlantic states, including DE, MD, PA, VA, and WV):

Apply 1.0 - 1.5 pts./A of **StreliaS II** + 0.5 - 1.0 pt./A of Banvel or Clarity® by ground equipment when pigweed plants are less than 3 inches tall and before corn exceeds 5 inches in height in a minimum of 20 gals. of spray per acre. Use the lower rate on *coarse-textured* and low organic matter soils. Use the higher rate on *fine-textured* and high organic matter soils.

Restriction:

1. Do not apply with aircraft.

Precaution:

1. Avoid drift to sensitive nontarget plants, such as soybeans, during application, or injury may occur.

TANK MIXTURE WITH AATREX OR PRINCEP + PROWL FOR PROLONGED CONTROL OF LAMBSQUARTERS AND PIGWEED IN FIELD CORN ONLY (NORTHEAST U.S., INCLUDING MI, IN, KY, AND STATES EAST OF THESE)

For prolonged control of lambsquarters and pigweed, in addition to a broad spectrum of annual broadleaf and grass weeds, **StreliaS II** in tank mix combination with AAtrex* or Princep + Prowl 4E may be applied after planting, but before corn or weeds emerge. Apply by ground equipment in a minimum of 10 gals. of water or 20 gals. of liquid fertilizer. Apply by air in a minimum of 5.0 gals. of water. Refer to **Table 3** of this label for rates of **StreliaS II**, AAtrex, or Princep to be applied. Apply Prowl 4E according to the rates in **Table 4**.

*Do not apply **StreliaS II** in tank mix combination with AAtrex 80W+ Prowl, as this combination is not compatible. Other AAtrex formulations may be used.

Mixing Instructions: See **Comment No. 1** following **Table 2**.

Table 4: Prowl 4E - Broadcast Rates Per Acre

Soil Texture	Percent Organic Matter in Soil		
	Less Than 1.5%	1.5 - 3%	Over 3%
COARSE	1.5 - 2.0 pts.	2.0 pts.	3.0 pts.
MEDIUM	2.0 pts.	3.0 pts.	3.0 pts.
FINE	2.0 pts.	3.0 pts.	3.0 pts.

Observe all directions for use, precautions, and limitations on the respective product labels when applying these products in tank mix combination. Refer to the Prowl 4E label for replanting instructions in the event of crop loss.

TANK MIXTURE WITH AATREX OR PRINCEP, OR AATREX + PRINCEP WITH GRAMOXONE BRANDS, LANDMASTER BW, TOUCHDOWN, OR ROUNDUP FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, state seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone brands, Landmaster BW, Touchdown brands, or Roundup brands should be tank mixed with **StreliaS II** + AAtrex, **StreliaS II** + Princep, or **StreliaS II** + AAtrex + Princep. See **Comment No. 7** following **Table 2**. The **StreliaS II**, **StreliaS II** + AAtrex or Princep, or **StreliaS II** + AAtrex + Princep portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for **StreliaS II**, **StreliaS II** + AAtrex or Princep, or **StreliaS II** + AAtrex + Princep - Preplant Surface, Preplant incorporated, or Preemergence.



Application:

Apply before, during, or after planting, but before the corn emerges. Add Gramoxone brands, Landmaster BW, Touchdown brands, or Roundup brands and apply as directed on the product label.

Gramoxone Brands:

Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

Restrictions: Do not apply combinations containing Gramoxone brands in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

Landmaster BW:

27 - 54 ozs./A depending on weed species and size. See the Landmaster BW label for weeds controlled, labeled rates for specific weeds, and other information concerning use.

Touchdown Brands or Roundup Brands:

See the Touchdown brand or Roundup brand label for weeds controlled, rates, and other use directions.

Apply in 20 - 60 gals. of water or fluid fertilizer per acre with ground equipment.

On *coarse soils*, apply 1.0 pt./A of **StreliaS II** with 1.3 lbs. of AAtrex Nine-0* or Princep Caliber 90*, or with 0.7 lb. of AAtrex Nine-0** + 0.7 lb. of Princep Caliber 90**. On *medium soils*, apply 1.33 pts./A of **StreliaS II** with 1.8 lbs. of AAtrex Nine-0 or Princep Caliber 90, or with 0.9 lb. of AAtrex Nine-0 + 0.9 lb. of Princep Caliber 90. On *fine soils****, apply 1.33 - 1.67 pts./A of **StreliaS II** with 1.8 - 2.2 lbs. of AAtrex Nine-0 or Princep Caliber 90, or with 0.9 - 1.1 lbs. of AAtrex Nine-0 + 0.9 - 1.1 lbs. of Princep Caliber 90.

* Use Princep in preference to AAtrex when heavy infestations of crabgrass or fall panicum are expected.

** When using the tank mixture of **StreliaS II** + AAtrex Nine-0 + Princep Caliber 90, use equal rates of AAtrex and Princep as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of AAtrex + Princep instead of the 1:1 ratio given. (Example: Total AAtrex Nine-0 + Princep Caliber 90 = 1.8 lbs./A, use 0.6 lb. of AAtrex + 1.2 lbs. of Princep, respectively.) Refer to **Comment No. 2** following **Table 2** for AAtrex 4L and Princep 4L conversions.

*** For cocklebur, yellow nutsedge, and velvetleaf control on *fine-textured soils* above 3% organic matter, apply 2.25 lbs./A of AAtrex Nine-0, or equivalent rate of AAtrex 4L, or the same total amount of AAtrex + Princep with 1.33 - 1.67 pts./A of **StreliaS II**.

TANK MIXTURE WITH AATREX; OR AATREX + 2,4-D; OR AATREX + 2,4-D + BANVEL FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, **StreliaS II** applied in combination with AAtrex will kill most emerged small annual weeds. Apply **StreliaS II** + AAtrex before, during, or after planting, but before corn emerges, according to the rates in **Table 3**.

Where heavy crop residues exist, add 0.8 - 1.6 pts./A of an appropriately labeled 3.8 lbs. a.i./gals. of 2,4-D amine (such as Weedar 64, Weedar 64A, DMA-4 Herbicide, Weedone® 638 or Formula 40) to the spray tank last and apply in a minimum of 25 gals. of carrier per acre.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds, and therefore, are recommended instead of water. Add X-77 surfactant at 1.0 - 2.0 qts./100 gals. of diluted spray, or another appropriate surfactant at its recommended rate, or add crop oil concentrate plus 28% liquid nitrogen (or equivalent). Apply before weeds exceed 3 inches in height. If alfalfa is present, add Banvel to the spray mixture at 0.33 - 0.5 pt./A and apply before alfalfa exceeds 6 inches in height.

For fields with existing sod grasses (e.g., bromegrass, orchardgrass, rye, or timothy), when existing weeds exceed 3 inches in height or when very dry conditions exist, add Gramoxone brands at the rate indicated on the product label in place of or in addition to 2,4-D, as indicated above. Do not apply Gramoxone brands in suspension-type liquid fertilizer. Observe all directions for use, precautions, and limitations on the respective product labels when applying these products in tank mix combination.

TANK MIXTURE WITH MARKSMAN IN CONSERVATION TILLAGE - FIELD AND SILAGE CORN

In conservation tillage systems where corn is planted directly into a cover crop or previous crop residue, **StreliaS II** + Marksman will kill most emerged small annual weeds. Apply **StreliaS II** + Marksman before, during, or after planting, but before corn emergence on *medium* and *fine soils* with greater than 2.5% organic matter. For fields with existing vegetation exceeding 3 inches in height or when very dry conditions exist, add Gramoxone brands at its standard rate. **StreliaS II** + Marksman may be applied postemergence to corn less than 3 inches tall and before weedy grasses exceed the 2-leaf stage.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds. Do not apply Gramoxone brands in suspension-type liquid fertilizer or use on emerged corn.

Refer to the Marksman label and follow all directions, limitations, precautions, and information regarding application and use in corn.

TANK MIXTURE WITH BALANCE PRO - FIELD CORN ONLY

StreliaS II and Balance Pro have a complementary response and weed control profile which allows various tank mix rate combinations to be considered. The addition of Balance Pro will improve the control of certain problem weeds, including Texas panicum, woolly cupgrass, and wild proso millet. **StreliaS II** improves both the duration and spectrum of annual grass and small-seeded broadleaf weed control, in particular foxtails (yellow foxtail) and witchgrass, and yellow nutsedge.

To reduce the risk of an adverse crop response, the Balance Pro label does not allow applications to *coarse-textured soils* with less than 1.5% organic matter and warns about applications to all soils with less than 1.5% organic matter or with pH greater than 7.5, as well as applications made to areas in fields with clay knolls, eroded hillsides, and exposed subsoil. **StreliaS II** has no adverse crop response warnings or use restrictions.

Listed below are compensating rate options for combinations of **StreliaS II** and Balance Pro, i.e., higher rates of **StreliaS II** are combined with lower rates of Balance Pro and vice versa. Select a rate option for **StreliaS II** plus Balance Pro by weighing the intensity of problem weed pressure (population presence and density) and your acceptance for risk of an adverse crop response. For example, where Texas panicum, woolly cupgrass, or wild proso millet is a primary target weed, use a tank-mix combination with a higher Balance Pro rate for the given soil type.

Where your acceptance of an adverse crop response risk is low and/or a more general weed spectrum is targeted (especially yellow foxtail, witchgrass, or yellow nutsedge), use a tank-mix combination with a higher **StreliaS II** rate for the given soil type. Where a target weed is listed as controlled on both product labels, a tank-mix combination option including intermediate rates of both products may be used. Where a target weed is listed as controlled on only one product label, do not apply a rate of that product below what is listed for that weed on the individual product label, or unacceptable control may result. Follow all other directions for use, rate limitations, precautions, and restrictions on both the **StreliaS II** and Balance Pro product labels.

For Coarse-textured Soils: Where 1.5 or 1.88 ozs./A of Balance Pro is used, 1.0 - 1.33 pts./A of **StreliaS II** may be applied. Do not use Balance Pro on *coarse-textured soils* with less than 1.5% organic matter.

For Medium-textured Soils: Where 1.5 ozs./A of Balance Pro is used, rates as low as 1.33 pts./A of **StreliaS II** may be applied. Where 1.88 or 2.25 ozs./A of Balance Pro is used, rates as low as 1.0 pt./A of **StreliaS II** may be applied. **StreliaS II** can be used in combinations with Balance Pro at rates up to 1.67 pts./A on *medium-textured soils*.

For Fine-textured Soils: Where 1.5 ozs./A of Balance Pro is used, rates as low as 1.33 pts./A of **StreliaS II** may be applied if the soil organic matter is less than 3%; if the soil organic matter is 3% or greater, 1.67 pts./A of **StreliaS II** should be applied. Where 1.88 or 2.25 ozs./A of Balance Pro is used, rates as low as 1.33 pts./A of **StreliaS II** may be applied. Where 3.0 ozs./A or more of Balance Pro is used, rates as low as 1.0 pt./A of **StreliaS II** may be applied. **StreliaS II** can be used in combinations with Balance Pro at rates up to 2.0 pts./A on *fine-textured soils* if the organic matter is 3% or greater.

TANK MIXTURES FOR POSTEMERGENCE SALVAGE WEED CONTROL IN FIELD CORN ONLY

For postemergence control of weeds in specific types of field corn, the **StreliaS II** combinations listed below may be used. Full season weed control from early preplant, preplant incorporated, or preemergence treatments can lead to maximum yield potential under competition-free conditions. However, if control of emerged weeds is needed, a postemergence program listed below can be applied to provide residual control for the remainder of the season.

Restrictions:

1. Follow all label directions, instructions, precautions, and limitations for each product used.
2. Do not use fluid fertilizer with these mixtures or corn injury may occur.
3. For each tank mixture with **StreliaS II**, apply only to the specific field corn type specified on the tank mix product label.



Precaution:

1. In row weed control may be reduced because of lack of coverage when applied to corn over 4 inches tall.

StreliaS II + Liberty Herbicide or Ignite® 280 SL Herbicide: Postemergence Use in LibertyLink® Corn or Corn Warranted by Bayer CropScience as Being Tolerant to Liberty Herbicide or Ignite 280 SL Herbicide

These tank mixtures can be applied postemergence to weeds and corn from seed designated as LibertyLink or corn warranted by Bayer CropScience as being tolerant to Liberty Herbicide or Ignite 280 SL Herbicide. Liberty provides postemergence control of a broad spectrum of grass and broadleaf weeds and the **StreliaS II** provides residual control of grasses and certain broadleaf weeds listed in the label section **STRELIUS II APPLIED ALONE - Weeds Controlled**. Refer to section **STRELIUS II APPLIED ALONE - Preplant Incorporated or Preemergence** and use the minimum rate per soil texture and organic matter classification for season-long residual control from this tank mix combination with Liberty Herbicide or Ignite 280 SL Herbicide. Refer to the Liberty Herbicide or Ignite 280 SL Herbicide label for the postemergence application rates according to weed species and their maximum height at the time of postemergence application. Where multiple weed species are present, use the highest rate listed to control the species and growth stages present.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the **StreliaS II** and Liberty Herbicide or Ignite 280 SL Herbicide labels. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

StreliaS II + Touchdown Brands or Roundup Brands for Postemergence Application to Glyphosate-tolerant Corn

The tank mixture of **StreliaS II** + Touchdown brands or Roundup brands can be applied postemergence to weeds and to corn designated as glyphosate-tolerant. Application may be applied postemergence to glyphosate-tolerant corn from emergence until corn reaches 30 inches tall or the V8 stage (8 leaves with collars), whichever comes first. This mixture will provide postemergence control of weed species on the Touchdown brand or Roundup brand label and residual control of weed species on the **StreliaS II** label. Use the minimum **StreliaS II** rate postemergence with Touchdown or Roundup in glyphosate-tolerant corn as specified in the **CORN (ALL TYPES) - STRELIUS II ALONE- PREPLANT INCORPORATED OR PREEMERGENCE** section of this label according to soil texture and organic matter. Refer to the Touchdown brand or Roundup brand label and follow appropriate use directions, application procedures, precautions, and limitations. Refer to the Touchdown brand or Roundup brand label for directions for control of problem species. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

StreliaS II + Touchdown Brands or Roundup Brands + AAtrex for Postemergence Application to Glyphosate-tolerant Corn

The tank mixture of **StreliaS II** + AAtrex + Touchdown brands or Roundup brands can be applied postemergence to weeds and to corn designated as glyphosate-tolerant. Application may be applied postemergence to glyphosate-tolerant corn from emergence up to 12 inches in height. This mixture will provide postemergence control of weed species on the Touchdown brand or Roundup brand label and residual control of weed species on the **StreliaS II** + AAtrex label. Use the minimum **StreliaS II** + AAtrex rate postemergence with Touchdown or Roundup in glyphosate-tolerant corn as specified in the **CORN - STRELIUS II COMBINATIONS - TANK MIXTURE WITH AATREX OR PRINCEP, OR AATREX + PRINCEP - PREPLANT SURFACE, PREPLANT INCORPORATED OR PREEMERGENCE** section and **Table 3** of this label according to soil texture and organic matter.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the **StreliaS II**, AAtrex, and Touchdown brand or Roundup brand labels for application to glyphosate-tolerant corn. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

COTTON

STRELIUS II ALONE

Application:

Apply **StreliaS II** preemergence only in Area 1* at the rate of 0.5 - 1.0 pt./A on sandy loams, 0.66 - 1.33 pts./A on *medium soils*, or 1.0 - 1.33 pts./A on *fine soils*. Apply **StreliaS II** preplant incorporated or preemergence in Area 2** at 1.0 pt./A on sandy loams, 1.0 - 1.33 pts./A on *medium soils*, or 1.33 pts./A on *fine soils*. Apply **StreliaS II** postemergence to cotton and preemergence to weeds at 0.5 - 1.33 pts./A, according to the state limitations in the follow **Postemergence** section.

* Area 1 = AR, KS, LA, MS, TN, and Bootheel of MO

** Area 2 = NM, OK, and TX

Fall Application for Italian Ryegrass Control:

StreliaS II may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply **StreliaS II** at 1.33 - 1.67 pts./A in the fall (September 1 - December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower **StreliaS II** rate for *coarse-textured soils* and the higher rate for *fine-textured soils*. A tillage operation may precede the application. Do not incorporate to a depth greater than 2 - 3 inches if tillage follows the application of **StreliaS II**. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone brands can be tank mixed with **StreliaS II** to control emerged ryegrass. Refer to the Gramoxone brands label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with **StreliaS II** for control or improved control of other weeds present at the time of application.

Restrictions:

1. Do not apply **StreliaS II** to frozen ground.
2. If a spring application is made, the combined total amount of **StreliaS II** applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for cotton (2.6 pts./A, depending on soil texture), or illegal residues may result.

Preplant Incorporated (NM, OK, and TX Only):

Apply to the soil and incorporate into the top inch of soil immediately before planting, at planting, or after planting, but before crop or weed emerge. Use a rolling cultivator or similar implement to uniformly incorporate not more than 1 inch deep. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. Where furrow irrigation is used, wet the top of the bed for best results. If the crop is to be planted on beds, apply and incorporate after bed formation. Cotton should be planted below the zone of incorporation; i.e., at least 1 inch on *fine soils* and 1.5 inches on *coarse* and *medium soils*. If incorporated prior to planting, use a planter that will result in a minimum of soil disturbance.

Restrictions: For best control of yellow nutsedge and suppression of seedling johnsongrass, apply **StreliaS II** preplant incorporated at the maximum rate for the soil texture, whether applied alone or mixed with Caparol 4L.

Preemergence:

Apply to the soil surface at planting or after planting, but before weeds or crop emerge.

Postemergence:

Apply **StreliaS II** broadcast over the top or directed to the soil surface according to the rate limitations listed below by state. Over-the-top postemergence application may be made not later than 100 days before harvest, and directed-postemergence application may be made not later than 80 days before harvest. Application before weeds emerge or after clean cultivation to remove existing weeds is necessary since **StreliaS II** will not control emerged weeds. **StreliaS II** postemergence may be applied over any previous registered herbicide treatment. In sprinkler-irrigated areas, sprinkler irrigate after application with 1/2 - 1 inch of water (1/2 inch on *coarse-textured soils* to 1 inch on *fine-textured soils*) to incorporate **StreliaS II**. In furrow-irrigated areas, apply **StreliaS II**, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In non-irrigated areas, if at least 1/2 inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of **StreliaS II**.

VA, NC, SC, GA, FL, and AL: Apply **StreliaS II** postemergence at 1.0 - 1.33 pts./A.

TN, AR, KS, MS, MO, and LA: Apply **StreliaS II** postemergence at 0.5 - 1.33 pts./A.

TX, OK, NM, AZ, CA, and Clay Soils in AR: Apply **StreliaS II** postemergence at 1.0 - 1.33 pts./A before August 1.

Multiple Applications: Where weed pressure is heavy, difficult-to-control species are expected, or re-infestation may occur, and a weed control program is used, multiple applications of **StreliaS II** are effective when used as part of the weed control program. Apply as a preplant incorporated or preemergence treatment and follow with an application postemergence to cotton before weeds emerge or after clean cultivation to remove existing weeds, since **StreliaS II** will not control emerged weeds. Apply **StreliaS II** postemergence over a previous preplant or preemergence **StreliaS II** application as shown in the following table.



State	Multiple Strelius II Applications to Cotton		
	Preplant Incorporated or Preemergence (pts./A)	+	Postemergence (pts./A)
MS, LA, TN, AR, KS, MO	0.5 - 1.33 (Preemergence only)	+	0.5 - 1.33
TX, OK, NM	1.0 - 1.33	+	1.0 - 1.33 before August 1
NC, VA	1.0 - 1.33 (Preemergence only)	+	1.0 - 1.33

In sprinkler-irrigated areas, sprinkler irrigate after application with 1/2 - 1 inch of water (1/2 inch on *coarse-textured soils* to 1 inch on *fine-textured soils*) to incorporate **Strelius II**. In furrow-irrigated areas, apply **Strelius II**, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In nonirrigated areas, if at least 1/2 inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of **Strelius II**.

Restrictions: For best control of yellow nutsedge and suppression of seedling johnsongrass, apply **Strelius II** preplant incorporated, preemergence, or postemergence to cotton and preemergence to weeds at the maximum rate for the soil texture, whether applied alone or in combinations. Do not apply more than a total of 2.0 pts./A on *coarse soils* or 2.6 pts./A of **Strelius II** on *medium* and *fine soils* during a growing season. These treatments may be applied over previous registered herbicide treatments.

Precautions: To avoid crop injury:

1. Do not apply **Strelius II** on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed;
2. To avoid concentration in the seed furrow, do not make broadcast applications of **Strelius II** to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow;
3. In furrow-planted cotton, to avoid concentration in the furrow and potential injury, do not apply **Strelius II** postemergence until after first "knifing" or cultivation to level soil surface;
4. Do not apply over the top in fluid fertilizer or any other adjuvant, surfactant, oil, or other pesticide not listed in the **COTTON** section of this label, or injury may occur;
5. Do not apply on Taloka silt loam;
6. Do not use in Gaines County, TX.

Restriction: Do not graze or feed forage or fodder from cotton to livestock, or illegal residues may result.

COTTON

STRELIUS II COMBINATIONS

TANK MIXTURE WITH CAPAROL 4L

Strelius II tank mixtures with Caparol 4L may be applied preplant incorporated or preemergence in water or fluid fertilizer. When fluid fertilizer is used as a carrier for **Strelius II**, either alone or in combination with Caparol 4L, mix only the amount that will be sprayed in one operation. These mixtures should not be allowed to stand without agitation. Only use water as a carrier for postemergence-directed application.

In addition to those weeds controlled by **Strelius II** alone, **Strelius II** + Caparol 4L, applied preplant incorporated or preemergence, also controls the following weeds: junglerice, wild oats, annual morningglory, groundcherry, hairy nightshade, lambsquarters, malva, mustard, prickly sida (teaweed), purslane, ragweed, and shallow germinating seedlings of cocklebur and coffeeweed. As a postemergence-directed application, Caparol 4L provides postemergence control and residual control of weeds on its label, while **Strelius II** provides residual control of weed species on its label. **Strelius II** will not control emerged weeds.

Preplant incorporated or Preemergence:

Apply **Strelius II** + Caparol 4L, either preplant incorporated or preemergence, using the appropriate rate from **Table 5**. Plant cotton below the zone of incorporation; i.e., at least 1.0 inch on *fine soils* and 1.5 inches on *coarse* and *medium soils*. If incorporated before planting, use a planter that will result in a minimum of soil disturbance.

Table 5: Strelius II + Caparol 4L - Cotton (NM, OK, TX)

Use Areas	Soil Texture	Broadcast Rates Per Acre	
		Strelius II	Caparol 4L
ALL	Sand, loamy sand	DO NOT USE	
OK, Blacklands and Gulf Coast of TX	Loams	0.8 - 1.33 pts.	2.4 pts.
	Clays	1.33 pts.	4.8 pts.
Rio Grande Valley of TX	Loams	0.8 - 1.33 pts.	3.2 pts.
	Clays	1.33 pts.	4.8 pts.
NM; High Plains, Rolling Plains, Edwards Plateau of TX; and Southwest TX	Sandy loam	0.8 - 1.0 pt.	1.6 pts.
	Loams	0.8 - 1.33 pts.	2.4 pts.
	Sandy clay loams	1.33 pts.	2.4 pts.
	Other clay soils	1.33 pts.	3.2 pts.

Postemergence Directed (AR, AZ, CA, LA, MS, NM, OK, TN, TX, and MO):

Strelius II may be tank mixed with Caparol 4L in water and applied postemergence directed in cotton for control of emerged weeds listed on the Caparol 4L label and residual preemergence control of weeds controlled by **Strelius II** and Caparol 4L, or application may be made after cultivation for residual preemergence control. These treatments may be applied over previous registered treatments, including **Strelius II**, provided the maximum label rate of any product is not exceeded. Do not apply over the top of cotton or injury may occur.

Apply **Strelius II** + Caparol 4L in a minimum of 20 gals. of spray volume per acre. Follow the directions, limitations, and precautions on the Caparol 4L label when Caparol 4L is applied as a postemergence-directed application. Refer to the directions, limitations, and precautions for use of **Strelius II** under the **COTTON - STRELIUS II ALONE - Postemergence** section.

Precautions:

To avoid concentration in the seed furrow, do not make broadcast applications of **Strelius II** + Caparol 4L to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.

To avoid crop injury:

1. Do not apply on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed;
2. Do not apply in cut areas of newly leveled fields, or in areas of excess salt;
3. Do not apply to glandless cotton varieties;
4. Do not apply on Taloka silt loam.
5. Do not use in Gaines County, TX.

Restrictions: Do not graze or feed forage or fodder from cotton to livestock, or illegal residues may result.

Refer to the Caparol 4L label for further instructions and limitations.



TANK MIXTURE WITH COTORAN 4L

Apply **StreliaS II** in tank mixture with Cotoran 4L preemergence for control of those weeds controlled by **StreliaS II** alone and those as listed on the Cotoran 4L label. This combination will also control spotted spurge, hyssop spurge, nodding spurge, and prostrate spurge. Apply to the soil surface at planting or after planting, but before weeds or crop emerge, using the appropriate rates from **Table 6**. Apply the tank mixture postemergence to cotton but preemergence to weeds, or apply postemergence to both cotton and broadleaf weeds for control of weeds on the Cotoran 4L label. Apply as a directed, semi-directed, or over-the-top spray. **StreliaS II** will not control emerged weeds but will provide preemergence control of species on its label. Where rate ranges are given for Cotoran 4L, use the higher rate when applying postemergence to weeds that are 2 inches or less. These treatments may be applied over previous registered treatments, including **StreliaS II**, provided the maximum label rate of any product is not exceeded.

Mixing Instructions: Incompatibility may occur when tank mixing **StreliaS II** and Cotoran 4L. To help overcome this condition, fill the spray tank 1/4 full with water or fluid fertilizer and start agitation, add the Cotoran 4L and allow it to become dispersed. Add X-77 at 0.5% volume/volume final spray (4.0 pts./100 gals.), then add the **StreliaS II** and finally the rest of the water or fluid fertilizer. Agitate during mixing and application to maintain a uniform suspension. Do not use fluid fertilizer as a carrier for postemergence applications.

Table 6: StreliaS II + Cotoran 4L - Cotton

Soil Texture	Broadcast Rates Per Acre		
	StreliaS II (pts.)		Cotoran 4L (pts.)
	Area 1*	Area 2**	
Sand, loamy sand	DO NOT USE		
Sandy loam	0.5 - 1.0	0.8 - 1.0	2.04
Loam, silt loam, silt	0.66 - 1.33	1.0 - 1.33	2.04 - 3.23
Fine soil	1.0 - 1.33	1.33	3.23 - 4.08

* Area 1 = AR, LA, MS, Bootheel of MO and TN

** Area 2 = Eastern OK, Gulf Coast, Rio Grande Valley, and Eastern TX

Precautions:

1. Do not apply **StreliaS II** + Cotoran on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed, or crop injury may occur.
2. To avoid concentration in the seed furrow, do not make broadcast applications of **StreliaS II** + Cotoran to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.
3. The use of Cotoran following the use of a systemic insecticide at planting may result in crop injury.
4. Do not use on Taloka silt loam or crop injury may occur.
5. Do not use in Gaines County, TX.

Refer to the Cotoran labels for further instructions, precautions, and limitations.

Restrictions: To avoid possible illegal residues, do not feed treated forage or gin trash to livestock, or graze treated areas.

TANK MIXTURE OF STRELIA S II OR STRELIA S II + COTORAN WITH GRAMOXONE BRANDS, TOUCHDOWN BRANDS, OR ROUNDUP BRANDS FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where cotton is planted directly into a cover crop, stale seedbed, or previous crop residues, the contact herbicides Gramoxone brands, Touchdown brands, or Roundup brands may be added to a tank mix of either **StreliaS II** or **StreliaS II** + Cotoran. When used as directed, the Gramoxone brands portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Touchdown or Roundup combinations will control emerged annual and perennial weeds when applied as directed on the Touchdown or Roundup label. The **StreliaS II** and **StreliaS II** + Cotoran portion of the tank mixture provides preemergence control of the weeds listed on this label in the **StreliaS II** and **StreliaS II** + Cotoran sections, respectively.

Refer to the label of each product used in combination and observe the planting details, information regarding application, geographical restrictions, and all other precautions and limitations. Refer to **Mixing Instructions** under **TANK MIXTURE WITH COTORAN 4L** section.

Application:

Apply before, during, or after planting, but before the cotton emerges. Apply **StreliaS II** at 0.8 - 1.0 pt./A on sandy loams, *medium*-, and *fine-textured* soils. Refer to **Table 6** for the Cotoran 4L rates.

Gramoxone Brands:

Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

Restrictions: Do not apply combinations containing Gramoxone brands in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

Touchdown Brands or Roundup Brands:

See the Touchdown or Roundup label for weeds controlled, rates, and other use directions.

Restrictions:

1. Do not apply **StreliaS II** + Cotoran 4L + Roundup in tank mixture because of compatibility problems.
2. Do not use in Gaines County, TX.
3. Apply in 20 - 60 gals. of water or fluid fertilizer per acre with ground equipment.

Precautions:

1. If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stand for several days, or where the seeding slit has not been properly closed.
2. Refer to the Cotoran labels and the **Tank Mixture with Cotoran 4L** section of this label for further instructions, precautions, and limitations.

TANK MIXTURE WITH MSMA, MSMA + CAPAROL, OR MSMA + COTORAN

StreliaS II may be tank mixed with MSMA in water and applied postemergence directed for control of emerged weeds listed on the MSMA product label and residual preemergence control of weeds controlled by **StreliaS II**. The addition of Caparol or Cotoran will add control of weed species on their respective labels.

Postemergence Directed (AL, AR, AZ, CA, FL, GA, LA, MS, NC, NM, OK, SC, TN, TX, VA, and Bootheel of MO):

Apply **StreliaS II** + MSMA postemergence directed to cotton at least 3 inches tall according to the directions, limitations, and precautions on the MSMA product label, as well as the directions, limitations, and precautions for use of **StreliaS II** in the section for **COTTON - STRELIA S II ALONE - Postemergence**. Do not apply after first cotton bloom. These treatments may be applied over previous registered treatments, including **StreliaS II**, provided the maximum label rate of any product is not exceeded. Cotoran or Caparol may be added to the **StreliaS II** + MSMA tank mixture according to the respective label directions for application to cotton at least 3 inches tall. When these mixtures are used, follow the mixing instructions for **StreliaS II** + Caparol or Cotoran and then add the MSMA product.

Do not use **StreliaS II** in tank mix with premixes of MSMA plus herbicides other than those registered for use in tank mixture with **StreliaS II** on cotton.



TANK MIXTURE WITH TREFLAN FOR POST-DIRECTED FOLLOWED BY SOIL INCORPORATION APPLICATIONS

StreliaS II may be applied as a tank mixture with Treflan in cotton for improved late-season weed control when used as an incorporated lay-by type application. This combination may be applied after the cotton is at least 3 inches tall and has reached the 4 true-leaf stage. Make the application directed to the soil surface and away from the crop foliage. Incorporate using a sweep or rolling type cultivator to provide uniform and shallow mixing into the top 2 inches of soil. Refer to each product label for the appropriate application rates by soil type and for this application timing, and follow all product use limitations and restrictions.

TANK MIXTURE WITH TOUCHDOWN BRANDS OR ROUNDUP BRANDS FOR USE ON GLYPHOSATE-RESISTANT COTTON ONLY

Apply **StreliaS II** as a tank mixture with Touchdown or Roundup in water postemergence over-the-top or postemergence directed for control of emerged weeds listed on the Touchdown or Roundup labels and for residual preemergence control of weeds listed on the **StreliaS II** label. See the **COTTON - STRELIAUS II ALONE - Postemergence** section of this label for rates and timings of **StreliaS II** and follow the Touchdown and Roundup label for their respective rates, application methods, and application timing restrictions. Do not add additional spray adjuvants, surfactants, fertilizer additives, or pesticides to this tank mixture if applied postemergence over-the-top or unacceptable injury may occur. Refer to the Touchdown brand or Roundup brand label and follow appropriate use directions, application procedures, precautions, and limitations.

Restrictions:

1. Do not apply this tank mixture postemergence to any cotton variety unless it is designated glyphosate-resistant and unless the Touchdown or Roundup formulation being used is registered for postemergence use in Glyphosate-resistant Cotton.
2. Do not apply Touchdown or Roundup postemergence over-the-top to cotton past the growth stage limit specified on their respective labels.
3. Do not use on sand or loamy sand soils in Gaines County, TX.

Precaution: Postemergence over-the-top applications of this tank mixture may cause temporary injury in the form of necrotic spotting to exposed cotton leaves, which will not affect normal plant development.

SOYBEAN

IMMATURE SEED

StreliaS II may be applied preplant or preemergence for the control or suppression of grass and small-seeded weeds in immature-seed soybean or other food-grade soybeans. For specific rates, see the rate table listed below.

Preplant Surface-applied:

For minimum-tillage or no-tillage systems only, **StreliaS II** alone may be applied up to 45 days before planting. Use only split applications for treatments made 30 - 45 days before planting, with 2/3 the listed broadcast rate for the crop and soil texture applied initially and the remaining 1/3 applied at planting. Treatments less than 30 days before planting may be made either as a split or a single application. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (e.g., Gramoxone brands, Touchdown, or Roundup). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated:

Apply **StreliaS II** to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a pre-plant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate **StreliaS II** after bed formation, unless specified otherwise.

Preemergence:

Apply **StreliaS II** during planting (behind the planter) or after planting, but before weeds emerge.

StreliaS II Broadcast Rates Per Acre

Soil Texture	Percent Organic Matter in Soil	
	< 3%	≥ 3%
MEDIUM	1.33 - 1.67 pts.	1.33 - 1.67 pts.
FINE	1.33 - 1.67 pts.	1.67 - 2.0 pts.

Precautions:

1. **StreliaS II** will not control emerged weeds.

Restrictions:

1. Do not cut for hay within 120 days following a **StreliaS II** application.
2. Do not use for forage within 60 days following a **StreliaS II** application.
3. Do not apply more than 2.0 pts./A of **StreliaS II** during any one crop year.

GRASSES GROWN FOR SEED (ID, OR, WA)

STRELIAUS II APPLIED ALONE

To control weeds and volunteer grasses in established grasses grown for seed, apply **StreliaS II** to established stands of tall fescue, orchard grass, perennial ryegrass, fine fescue, bentgrass, and Kentucky bluegrass just before, during, or immediately following the first fall rains or just before or during a late summer or early fall irrigation, but before target grasses emerge. The seed crop must have had one seed harvest or been established at least one year. The post-harvest residue (straw) should be evenly spread, removed, or burned before applying **StreliaS II**. Rainfall or irrigation is required after application and before weed, emergence for best control. **StreliaS II** will provide preemergence control/suppression of volunteer seedlings of perennial ryegrass, fine fescue species, tall fescue, orchardgrass, bentgrass, and Kentucky bluegrass. **StreliaS II** will control those weed species listed in the **STRELIAUS II APPLIED ALONE** section of the **StreliaS II** label and will suppress or control rattail fescue, annual bluegrass, Italian ryegrass, California brome, downy brome, and roughstalk bluegrass.

Apply **StreliaS II** by ground equipment in a minimum of 10 gallons of water per acre using the rate listed below according to grass species.

Established Grass Crop Grown for Seed	pts./A
Fine fescue species	1.0
Perennial ryegrass	1.0
Bentgrass	1.0 - 1.33
Kentucky bluegrass	1.0 - 1.33
Orchardgrass	1.0 - 1.33
Tall fescue	1.0 - 1.33



Precautions:

1. Tank mixtures with other pesticides, or the addition of an adjuvant, can increase the risk of crop injury.
2. Application to perennial ryegrass and fine fescue stands under stress may cause crop injury.
3. If weed escapes occur following **StreliaS II** application, an application of a postemergence herbicide may be necessary to control escapes. When making such an application, follow all directions, precautions, and limitations on the label of the postemergence herbicide.
4. Control may be decreased if excessive straw from the previous harvest is present at application and/or insufficient rainfall/irrigation occurs.

Restrictions:

1. Do not apply **StreliaS II** more than once per crop year.
2. Do not apply after November 15 or poor control may result.

To avoid possible illegal residues:

1. Do not graze forage regrowth for 60 days following application west of the Cascades.
2. In areas east of the Cascades, do not graze forage regrowth for 150 days following application.
3. Hay may be harvested anytime between seed harvest and the next application of S-metolachlor.

HORSERADISH

Apply a single application of **StreliaS II** at a broadcast rate of 1.0 - 1.33 pts./A to the soil surface after planting, but before weed or crop emergence (i.e., preemergence). Use lower rates on soils relatively *coarse-textured* and higher rates on *fine-textured* soils. A band application may also be used, applying proportionally less spray mixture on the area actually treated. **StreliaS II** will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical means.

Restrictions:

1. Make only one application of **StreliaS II** per crop.
2. Do not apply more than 1.33 pts./A of **StreliaS II** per crop.
3. Harvest horseradish at normal timing.

PEANUTS

STRELIUS II ALONE

Apply **StreliaS II**, either preplant incorporated, postplant incorporated, or preemergence, using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of **StreliaS II** alone under **APPLICATION PROCEDURES. Postplant Incorporated:** Apply and shallowly incorporate **StreliaS II** into the soil after planting, but before peanut germination. Incorporation depth and incorporating implements must be kept above the seed, or seed will be damaged.

Apply **StreliaS II** alone, preplant incorporated, postplant incorporated, or preemergence, at a broadcast rate of 1.0 - 1.33 pts./A in the Southeast* and 0.8 - 1.33 pts./A in NM, OK, and TX.

* In the Southeast, use 1.33 - 2.0 pts./A and apply preemergence for partial control of Florida beggarweed.

Restrictions:

1. **StreliaS II** alone may be applied as directed after any of the following preplant incorporated herbicides when used according to their label directions: Treflan E.C. at 1.0 pt./A; Sonalan at 1.25 - 3.0 pts./A; Pursuit at 0.25 pt./A; or Prowl at 1.0 - 2.0 pts./A.
2. Do not graze or feed peanut forage or fodder to livestock for 30 days following application.
3. Do not apply within 90 days of harvest, or illegal residues may result.

PEANUTS

STRELIUS II COMBINATIONS

TANK MIXTURE OR SEQUENTIALLY WITH PURSUIT

The tank mixture or sequential treatment of **StreliaS II** and Pursuit controls all weeds controlled by **StreliaS II** alone and by Pursuit alone. Refer to the **STRELIUS II APPLIED ALONE** section for weeds controlled by **StreliaS II** and to the Pursuit label for weeds controlled by Pursuit.

Refer to the respective labels for application methods, timing, rates, restrictions, and precautions; and use in accordance with the more restrictive label. Do not exceed the label rate of either product **StreliaS II** will not control emerged weeds.

TANK MIXTURE WITH SONALAN

The tank mixture controls all weeds controlled by **StreliaS II** alone and by Sonalan alone. Refer to the **STRELIUS II APPLIED ALONE** section for weeds controlled by **StreliaS II** and to the Sonalan label for weeds controlled by Sonalan.

Apply **StreliaS II** + Sonalan preplant incorporated, using the appropriate rate from **Table 7**. Follow recommended soil preparation procedures for Sonalan.

Table 7: StreliaS II + Sonalan - Peanuts

Soil Texture	Broadcast Rates Per Acre			
	Southeast		NM, OK, TX	
	StreliaS II	Sonalan	StreliaS II	Sonalan
COARSE	1.0 - 1.33 pts.	1.25 - 2.0 pts.	0.8 - 1.33 pts.	1.25 - 2.0 pts.
MEDIUM	1.0 - 1.33 pts.	1.75 - 2.5 pts.	0.8 - 1.33 pts.	1.75 - 2.5 pts.
FINE	1.0 - 1.33 pts.	2.25 - 3.0 pts.	0.8 - 1.33 pts.	2.25 - 3.0 pts.

Restrictions: Follow all use directions, limitations, precautions, and information regarding application to peanuts on the **StreliaS II** and Sonalan labels.

TANK MIXTURE WITH PROWL

StreliaS II + Prowl applied preplant incorporated controls all weeds controlled by **StreliaS II** alone plus Texas panicum, field sandbur, johnsongrass from seed, lambsquarters, kochia, annual spurge, and other species on the Prowl label. Apply **StreliaS II** + Prowl by ground or by aerial equipment within 14 days before planting. Incorporate into the top 1 - 2 inches of soil before planting and within 7 days of application, using a finishing disk or similar implement capable of providing uniform incorporation. If peanuts will be planted on beds, apply and incorporate after bed formation. Refer to the incorporation instructions of the respective labels for additional directions.



Apply **StreliaS II** + Prowl preplant incorporated, using the appropriate rates from **Table 8**.

Table 8: StreliaS II + Prowl - Peanuts

Soil Texture	Broadcast Rates Per Acre	
	NM, OK, TX	Other Peanut Growing States
	StreliaS II + Prowl	StreliaS II + Prowl
Sand, loamy sand	0.8 + 1.0 - 1.5 pts.	1.0 - 1.33 + 1.5 - 2.0 pts.
Sandy loam	0.8 - 1.0 - 1.5 pts.	1.0 - 1.33 + 1.5 - 2.0 pts.
Fine soil	1.33 + 1.0 - 1.5 pts.	1.33 + 1.5 - 2.0 pts.

Restrictions: Follow all use directions, limitations, precautions, and information regarding application to peanuts on the **StreliaS II** and Prowl labels.

TANK MIXTURE WITH GRAMOXONE BRANDS

StreliaS II + Gramoxone brands applied at ground craking will control or suppress small (1- to 6-inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **STRELIA S II APPLIED ALONE** section of this label. Apply Gramoxone brands plus the appropriate **StreliaS II** rate from the **PEANUTS - STRELIA S II ALONE** section in a minimum spray volume of 20 gals./A with ground equipment. Refer to the Gramoxone brands label and follow all directions, limitations, and restrictions.

TANK MIXTURE WITH GRAMOXONE BRANDS + BASAGRAN

The addition of Basagran to the **StreliaS II** + Gramoxone brands mixture will result in improved control of such problem broadleaf weeds as prickly sida, cocklebur, smartweed, and bristly starbur. **StreliaS II** + Gramoxone brands + Basagran applied at ground craking will control or suppress small (1- to 6-inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **STRELIA S II APPLIED ALONE** section of this label. Apply Basagran + Gramoxone brands with the appropriate **StreliaS II** rate from the **PEANUTS - STRELIA S II ALONE** section in a minimum spray volume of 20 gals./A with ground equipment. Refer to the Gramoxone brands and Basagran labels and follow all directions, limitations, and restrictions.

TANK MIXTURE WITH GRAMOXONE BRANDS + BUTYRAC 200

The addition of Butyrac 200 to the **StreliaS II** + Gramoxone brands mixture will result in improved control of such problem broadleaf weeds as sicklepod, morning glory, and cocklebur. **StreliaS II** + Gramoxone brands + Butyrac 200 applied at ground craking will control or suppress small (1- to 6-inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **STRELIA S II APPLIED ALONE** section of this label. Apply Gramoxone brands + Butyrac 200 with the appropriated **StreliaS II** rate from the **PEANUTS STRELIA S II ALONE** section in a minimum spray volume of 20 gals./A with ground equipment. Refer to the Gramoxone brands and Butyrac 200 labels and follow all directions, limitations, and restrictions.

TANK MIXTURE WITH BASAGRAN

StreliaS II + Basagran applied at ground craking will control species on the Basagran label and provide residual control of species listed in the **STRELIA S II APPLIED ALONE** section of this label. Apply 1.0 - 2.0 pts./A of Basagran in 20 gals./A, depending on weed species and stage of growth as specified on the Basagran label, with the appropriate **StreliaS II** rate from the **PEANUTS - STRELIA S II ALONE** section. A second Basagran application may be made in all peanut-growing areas, if needed. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

TANK MIXTURE WITH BASAGRAN + BUTYRAC 200

StreliaS II + Basagran + Butyrac 200 applied at ground craking will control species on the Basagran label and on the Butyrac 200 label, especially morning glories. Apply 1.5 - 2.0 pts./A of Basagran + 8.0 fl. ozs./A of Butyrac 200 in 20 gals./A, depending on weed species and stage of growth as specified on the Basagran label, with the appropriate **StreliaS II** rate from the **PEANUTS - STRELIA S II ALONE** section. A second Basagran + Butyrac 200 application may be made in all peanut-growing areas, if needed. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

SEQUENTIALLY WITH STORM®

Apply **StreliaS II** according to the directions for **StreliaS II Alone** and follow with a postemergence treatment of Storm as specified on its label for the control of weeds listed on the **StreliaS II** label and on the Storm label. Refer to the **PEANUTS - STRELIA S II ALONE** section and to the Storm label and follow all directions, limitations, and restrictions for each product.

MULTIPLE APPLICATIONS

Where weed pressure is heavy or where species difficult to control are expected, **StreliaS II**, is most effective when used as follows:

Southeast Only (AL, FL, GA, NC, SC, VA)

Preplant Incorporated:

Apply **StreliaS II** preplant incorporated as directed under **PEANUTS - STRELIA S II ALONE** or apply **StreliaS II** + Balan preplant incorporated as directed previously in this section. Refer to the respective section for weeds controlled.

OR

Preemergence to Before "Ground Cracking":

Apply **StreliaS II** any time from preemergence to before "ground cracking" at 1.0 - 2.0 pts./A for extended control of weeds not yet emerged. **Do not use StreliaS II after peanut emergence.** If peanuts have emerged, use an unsafened S-metolachlor product according to its label: **Peanuts - Combinations - Multiple Applications.**

Follow the PPI or PRE Application By:

Lay-by:

Do not use StreliaS II. Apply an unsafened S-metolachlor product at lay-by as directed under the **Peanuts - Alone** section of the unsafened S-metolachlor product label.

Restrictions:

1. Do not apply more than the equivalent of 2.67 lbs. of active ingredient of **StreliaS II** per acre during any one year, or illegal residues may result. If an unsafened S-metolachlor product is used as a sequential treatment, the lbs. of active ingredient (1.0 pt. = 0.95 lb.) plus the lbs. of active ingredient of **StreliaS II** should not exceed 2.67 lbs. **Do not use StreliaS II after peanuts have emerged.**
2. Do not graze or feed peanut forage or fodder to livestock for 30 days following application.
3. Do not apply within 90 days of harvest, or illegal residues may result.

Southwest Only (NM, OK, TX)

1st Application:

Apply **StreliaS II** preplant incorporated or preemergence to before "ground cracking" as directed under **PEANUTS - STRELIA S II ALONE** or apply **StreliaS II** + Balan preplant incorporated as directed previously in this section. **Do not use StreliaS II after peanut emergence.** If peanuts have emerged, use an unsafened S-metolachlor product according to its label.

2nd Application:

Do not use StreliaS II. Apply an unsafened S-metolachlor product at lay-by as directed under the **Peanuts - Alone** section of the unsafened S-metolachlor product label. Use only when late germinating weeds are expected to be a problem. Refer to the product **Applied Alone** section for a list of weeds controlled.



Restrictions:

1. Do not apply more than the equivalent of 2.67 lbs. of active ingredient of **Strelius II** per acre during any one year, or illegal residues may result. If an unsafened *S*-metolachlor product is used as a sequential treatment, the lbs. of active ingredient (1.0 pt. = 0.95 lb.) plus the lbs. of active ingredient of **Strelius II** should not exceed 2.67 lbs. **Do not use Strelius II after peanuts have emerged.**
2. Do not graze or feed peanut forage or fodder to livestock for 30 days following application.
3. Do not apply within 90 days of harvest, or illegal residues may result.

BEANS, PEAS, AND LENTILS**STRELIUS II ALONE**

Beans, peas, and lentils, including garbanzo, great northern beans, kidney beans, lima beans, mung beans, navy beans, peas (English*; southern peas, such as blackeye, pinkeye, crowder, etc.), pinto beans, snap beans (green, wax, string), lentils, and lupines (sweet, white, white sweet, and grain).

Fall Application:

1. Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
2. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
3. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 - 2.0 pts./A on *medium-textured* and 2.0 pts./A on *fine-textured* soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2 - 3 inches. Minimize furrow and ridge formation in the tillage operations.

Restrictions: If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for beans, peas, and lentils or illegal residues may result.

Spring Application:

Apply **Strelius II**, either preplant incorporated or preemergence, using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of **Strelius II** alone under **APPLICATION PROCEDURES**. On *coarse soils* with less than 3% organic matter, apply 1.0 - 1.33 pts./A of **Strelius II** or 1.33 pts./A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts./A of **Strelius II**. On *fine soils*, apply 1.33 - 1.67 pts./A of **Strelius II** if organic matter content is less than 3%, or 1.67 - 2.0 pts./A if organic matter content is 3% or greater.

* On English peas, use only preemergence applications. If soils are cold and wet during pea germination and emergence, the use of **Strelius II** may delay maturity and/or reduce yields.

Restrictions: To avoid possible illegal residues,

1. Do not cut for hay within 120 days following a **Strelius II** application.
2. Do not use for forage within 60 days following a **Strelius II** application.
3. Do not apply more than 2.0 pts./A of **Strelius II** during any one crop year.

BEANS, PEAS, AND LENTILS**STRELIUS II COMBINATIONS**

Restrictions: When applying **Strelius II** in combination on beans, peas, and lentils, do not cut for hay within 120 days following application, or illegal residues may result.

TANK MIXTURE AND SEQUENTIAL APPLICATIONS WITH EPTAM - BEANS (GREEN OR DRY)

This mixture controls all weeds controlled by **Strelius II** alone and by Eptam alone. Refer to the **STRELIUS II APPLIED ALONE** section of this label for weeds controlled by **Strelius II** alone and to the Eptam label for weeds controlled by Eptam.

Preplant Incorporated:

Follow instructions for use of **Strelius II** alone under **APPLICATION PROCEDURES**. **Sequential:** Apply Eptam alone preplant incorporated, as specified on that label. Follow with a preemergence application of **Strelius II**, at rates specified for **Strelius II** alone, during planting (behind the planter) or after planting, but before the weeds or crop emerge.

Refer to the **PRODUCT INFORMATION** section of this label and to the Eptam label for weather, cultural practices, and all other precautions and limitations that affect performance of these products.

Apply 2.5 - 4.5 pts./A of Eptam 7E* with **Strelius II** as specified. On *coarse soils*, apply 0.8 pt./A of **Strelius II** if organic matter content is less than 3%, or 1.0 pt./A if organic matter content is 3% or greater. On *medium soils*, apply 1.0 pt./A of **Strelius II** if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *fine soils*, apply 1.33 pts./A of **Strelius II** if organic matter is less than 3%, or 1.33 - 1.67 pts./A if organic matter is 3% or greater.

* Refer to the Eptam label for rate limitations depending on geographical area, and for species and varietal restrictions.

Restriction: Do not exceed 3.5 pts./A of Eptam 7E on small white beans or green beans grown on coarse-textured soils.

TANK MIXTURE WITH TREFLAN - BEANS (DRY - KIDNEY, NAVY, PINTO, ETC.; LIMA; AND SNAP)

Strelius II + Treflan tank mix applied preplant incorporated controls those weeds listed under **STRELIUS II APPLIED ALONE** and those weeds listed for Treflan alone on the Treflan label. **Strelius II** + Treflan may be applied by ground or by aerial equipment and incorporated up to 14 days prior to planting. Follow the procedures on this label and on the respective Treflan label using equipment that provides uniform 2-inch incorporation.

Apply **Strelius II** + Treflan tank mix using the appropriate **Strelius II** rate specified for **Strelius II** alone, and the Treflan rate from the Dry Beans, and the Lima and Snap Beans sections of the respective Treflan label. Choose the product rate for the specific soil texture/organic matter classification and weed species expected.

Restrictions: Follow all restrictions and precautions on the respective Treflan label and in the **BEANS, PEAS, AND LENTILS - STRELIUS II ALONE** section of this label.

POTATOES**STRELIUS II ALONE**

Apply **Strelius II**, either incorporated, preemergence, or postemergence to potatoes after hilling/lay-by, according to directions specified below for control of weeds listed under the **PRODUCT INFORMATION** section. Within a rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. For application by center pivot irrigation, see the **CENTER PIVOT IRRIGATION APPLICATION** section of this label.

Incorporated:

Apply **Strelius II** at 1.0 - 2.0 pts./A to the soil and incorporate into the top 3 inches before planting, using a finishing disk, harrow, rolling cultivator, or similar implement. Planting and later cultural practices should not bring untreated soil to the surface. Postplant incorporated application may be made any time after planting to drag-off, but before potato emergence. Use an implement that evenly distributes **Strelius II** in the top 2 inches of soil. Do not damage potato seed pieces or sprouts with incorporation equipment.



Preemergence:

Apply **Strelius II** at 1.0 - 2.0 pts./A, either after planting as a preemergence, delayed preemergence, after drag-off or hilling treatment, but before weeds emerge. Up to 2.5 pts./A of **Strelius II** alone may be used where soil organic matter is between 6% and 20%.

Postemergence After Hilling/Lay-by:

Apply 1.67 pts./A of **Strelius II** postemergence to potatoes through after hilling/at lay-by to control **Strelius II** sensitive species for remainder of the growing season. This application will not control emerged weeds. It may be applied over a previous **Strelius II** application, but do not apply more than 3.6 pts./A of **Strelius II** in a single crop season.

Precautions:

1. Do not use on muck or peat soils. If cool, wet soil conditions occur after application, **Strelius II** may delay maturity and/or reduce yield of Superior and other early maturing potato varieties.
2. These directions for use do not apply to sweet potatoes or yams.
3. Do not apply both as a preemergence and an incorporated treatment.

Restrictions: Potatoes treated with **Strelius II** should not be harvested within 60 days after the at-planting to drag-off application, or illegal residues may result.

POTATOES**STRELIUS II COMBINATIONS****TANK MIXTURE WITH SENCOR**

In addition to those weeds controlled by **Strelius II** alone, **Strelius II** applied in tank mix combination with, or sequentially with, any of the registered Sencor formulations also controls the following broadleaf weeds: cocklebur*, hairy nightshade*, hemp sesbania, jimsonweed*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard.

* Partially controlled.

Strelius II at 1.0 - 2.0 pts./A plus the labeled Sencor use rate may be used preemergence or postemergence to potatoes through after last hilling. Apply 1.0 - 1.33 pts./A of **Strelius II** on coarse soils and 1.33 - 2.0 pts./A on other soil textures. Within this rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. **Strelius II** will not control emerged weeds.

Refer to the Sencor label for precautionary statements, restrictions, application information, center pivot irrigation application, weeds controlled, and varietal limitations.

Precautions:

1. Postemergence applications to potatoes, except center pivot, should be made only as a directed or semi-directed spray to avoid chlorosis, minor necrosis, or leaf distortion.
2. These directions for use do not apply to sweet potatoes or yams.
3. Do not use this tank mixture on muck or peat soils.

Restrictions:

1. Potatoes treated with **Strelius II** in tank mixture with Sencor cannot be harvested within 60 days after application, or illegal residues may result.
2. Potatoes may not be harvested within 40 days after a lay-by application of **Strelius II**, or illegal residues may result.

STRELIUS II + LOROX DF TANK MIXTURE (EAST OF ROCKY MOUNTAINS)

Strelius II may be applied in a tank mix combination with any of the registered Lorox formulations as a preemergence broadcast application to potatoes. Apply to the soil surface after planting and before emergence of the crop or after final drag-off according to the rates specified in **Table 9**.

Table 9: Strelius II + Lorox DF - Potatoes (East of Rocky Mountains)

Soil Texture	Broadcast Rates Per Acre			
	1% to Less Than 3% Organic Matter		3 to 5% Organic Matter	
	Strelius II	Lorox DF	Strelius II	Lorox DF
COARSE Sandy loam	1.0 pt.	1.0 - 1.5 lbs.	1.33 pts.	1.5 - 2.0 lbs.
MEDIUM Loam, silt loam, silt	1.33 pts.	1.5 - 2.0 lbs.	1.67 - 2.0 pts.	2.0 - 2.5 lbs.

Precautions: To avoid crop injury,

1. Do not use on sands or loamy sand.
2. Do not incorporate or spray over the top of emerged potatoes.

Refer to the **PRODUCT INFORMATION** section of this label and to the Lorox label for precautionary statements, restrictions, application information, and weeds controlled.

TANK MIXTURE WITH PROWL 4E

In addition to the weeds controlled by **Strelius II** alone, this tank mixture with Prowl 4E controls such problem species as kochia, lambsquarters, purslane, annual spurge, stinging nettle, and others specified on the Prowl 4E label. Apply **Strelius II** + Prowl 4E preemergence, preemergence incorporated, or early postemergence, according to the specific directions on the Prowl 4E label, using the rates in **Table 10**.

Table 10: Strelius II + Prowl 4E - Potatoes

Soil Texture	Broadcast Rates Per Acre	
	Less Than 3% Organic Matter	More Than 3% Organic Matter
	Strelius II + Prowl 4E*	Strelius II + Prowl 4E*
COARSE	1.0 - 1.33 pts. + 1.0 - 1.5 pts.	1.0 - 1.33 pts. + 1.0 - 1.5 pts.
MEDIUM	1.33 pts. + 1.5 - 2.0 pts.	1.33 - 1.67 pts. + 2.0 - 3.0 pts.
FINE	1.33 - 1.67 pts. + 2.0 - 3.0 pts.	1.67 - 2.0 pts. + 3.0 pts.

* When using other formulations of Prowl, use equivalent rates of active ingredient.

Refer to the **Strelius II** and Prowl 4E labels and observe all directions, timings, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.



TANK MIXTURE WITH PROWL 4E + EPTAM

In addition to the weeds controlled by **StreliuS II** alone, this tank mixture will control those species on the Prowl 4E and Eptam labels. Refer to the **StreliuS II + Prowl 4E** labels for rates of those products and add Eptam 7E at 3.5 - 7.0 pts./A, depending on geographical area. Refer to the respective **StreliuS II**, Prowl 4E, and Eptam labels and observe all directions, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

PUMPKIN

STRELIUS II ALONE

Preemergence

Apply **StreliuS II** preemergence (before the weeds have emerged) at 1.0 to 1.33 pts./A as an inter-row or inter-hill application in pumpkin. Leave 1 foot of untreated area over the row, or 6 inches to each side of the planted hill and/or any emerged pumpkin foliage (inter-row or inter-hill means not directly over the planted seed or young pumpkin plants). Use the lower **StreliuS II** rate on soils light in texture (loamy sand or lighter) and low in soil organic matter (less than 3%). **StreliuS II** applied as a broadcast spray over the planted row or hill, or applications made directly to crop foliage, will increase the risk of injury (e.g., stand loss, delayed maturity, and loss of yield) to the pumpkin crop. Do not apply **StreliuS II** closer than 30 days before pumpkin harvest.

StreliuS II will not control emerged weeds, and thus should be applied before the weeds emerge. Weeds that are present should be controlled by another means, e.g., by mechanical means or by another herbicide.

RHUBARB

STRELIUS II ALONE

Apply **StreliuS II** at a broadcast rate of 0.67 - 1.33 pts./A to the soil surface in early spring, prior to crop emergence. Use lower rates on soils relatively coarse-textured and higher rates on *fine-textured soils*. A band application may also be used, applying proportionally less spray mixture on the area actually treated. **StreliuS II** will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

Restrictions:

1. Make only one application of **StreliuS II** per crop.
2. Do not apply more than 1.33 pts./A of **StreliuS II** per crop.
3. Do not harvest rhubarb within 62 days of the **StreliuS II** application.

SAFFLOWERS

STRELIUS II ALONE

Preplant Incorporated or Preemergence:

Follow instructions for use of **StreliuS II** alone under **APPLICATION PROCEDURES**.

On *coarse soils*, apply 1.0 - 1.33 pts./A of **StreliuS II** if organic matter content is less than 3%, or 1.33 pts./A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts./A of **StreliuS II**. On *fine soils*, apply 1.33 - 1.67 pts./A of **StreliuS II** if organic matter content is less than 3%, or 1.67 - 2.0 pts./A if organic matter content is 3% or greater.

GRAIN OR FORAGE SORGHUM (SEED TREATED WITH CONCEP®)

STRELIUS II ALONE

Apply **StreliuS II**, either preplant surface, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Apply **StreliuS II** alone only when the sorghum seed has been properly treated with Concep seed treatment. Preplant or preemergence applications of **StreliuS II** not treated with Concep seed treatment will result in crop death.

Fall Application for Italian Ryegrass Control:

StreliuS II may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply **StreliuS II** at 1.33 - 1.67 pts./A in the fall (September 1 - December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower **StreliuS II** rate for *coarse-textured soils* and the higher rate for *fine-textured soils*. A tillage operation may precede the application. Do not incorporate to a depth greater than 2 - 3 inches if tillage follows the application of **StreliuS II**. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone brands can be tank mixed with **StreliuS II** to control emerged ryegrass. Refer to the Gramoxone brands label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with **StreliuS II** for control or improved control of other weeds present at the time of application.

Restrictions:

1. Do not apply **StreliuS II** to frozen ground.
2. If a spring application is made, do not apply **StreliuS II** or any other product containing S-metolachlor the following spring to grain or forage sorghum, or illegal residues may result.

Preplant Surface-applied:

Refer to instructions for use of **STRELIUS II** under **APPLICATION PROCEDURES** section on this label. For minimum-tillage or no-tillage systems only, apply **StreliuS II** up to 45 days before planting in CO, IA, IL, KS, MO, NE, and SD. Use only split applications for treatments made 30 - 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 1.5 pts./A of **StreliuS II** on *medium soils* or 1.67 pts./A on *fine soils*. Treatments less than 30 days prior to planting may be made either as a split or single application. Apply 1.33 pts./A of **StreliuS II** on *coarse soils* not more than 2 weeks prior to planting. Under dry conditions, irrigation after application is recommended to move **StreliuS II** into the soil.

Preplant Incorporated or Preemergence:

Refer to instructions for use of **STRELIUS II** under **APPLICATION PROCEDURES** section on this label. Broadcast 1.0 - 1.33 pts./A of **StreliuS II** on *coarse soils*, 1.33 - 1.5 pts./A on *medium soils*, or 1.33 - 1.67 pts./A on *fine soils*.

Postemergence: Refer to instructions for use of **STRELIUS II** under **APPLICATION PROCEDURES** section on this label. **StreliuS II** may be applied broadcast postemergence at 1.0 - 1.33 pts./A on *coarse soils*, 1.33 - 1.5 pts./A on *medium soils*, or 1.33 - 1.67 pts./A on *fine soils*. **StreliuS II** will not control emerged weeds. Therefore, emerged weeds must be controlled by cultural or chemical means. When applied alone, **StreliuS II** will be safe to emerged sorghum. The risk of sorghum injury increases when adjuvants (e.g., non-ionic, crop oil), nitrogen sources (e.g., AMS, UAN), or fertilizers are applied with **StreliuS II**.

Precautions:

1. If sorghum seed is not properly treated with Concep seed treatment, preplant and preemergence applications of **StreliuS II** will severely injure the crop.
2. Under high soil moisture conditions prior to sorghum emergence, injury may occur following preplant and preemergence application of **StreliuS II**. The crop will normally outgrow this effect.
3. Do not use **StreliuS II** on sorghum grown under dry mulch tillage, or injury may occur.

Restrictions:

1. Except for the split preplant surface treatment, do not make more than one application per year, or illegal residues may result.
2. Do not apply **StreliuS II** postemergence within 75 days of harvest.



GRAIN OR FORAGE SORGHUM (SEED TREATED WITH CONCEP)

STRELIUS II TANK MIXTURES

StreliaS II preplant or preemergence (prior to sorghum emergence) tank mixtures with AAtrex may be applied in water or fluid fertilizer. Apply **StreliaS II** preplant or preemergence in tank mixtures only when the sorghum seed has been properly treated with Concep seed treatment. Preplant or preemergence applications of **StreliaS II** to sorghum not treated with Concep seed treatment will result in crop death.

IMPORTANT: FOR TANK MIXTURES WITH AATREX (OR OTHER BRANDS OF ATRAZINE) - If applying StreliaS II in tank mixture with AAtrex, all the restrictions and rate limitations on the AAtrex label must be followed if more restrictive/protective than those on this label. In addition, if AAtrex is/must be applied at rates lower than those listed on this label, broadleaf weed control may be affected. Refer to the AAtrex label for weeds controlled at the reduced rates.

Precautions:

1. Applications of **StreliaS II** + AAtrex on highly alkaline soils or on eroded areas where calcareous subsoils are exposed may cause sorghum injury.
2. If sorghum seed is not properly treated with Concep, **StreliaS II** + AAtrex may severely injure the crop.
3. Under high soil moisture conditions prior to sorghum emergence, injury may occur following preplant and preemergence application of **StreliaS II** + AAtrex. The crop will normally outgrow this effect.
4. Do not use **StreliaS II** + AAtrex on sorghum grown under dry mulch tillage, or injury may occur.

Restriction:

1. Except for the split preplant surface treatment, do not make more than one application per year, or illegal residues may result.

TANK MIXTURE WITH AATREX

In addition to the weeds controlled by **StreliaS II** alone, **StreliaS II** + AAtrex also controls the following broadleaf weeds when applied either preplant surface, preplant incorporated, or preemergence: cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

Preplant Surface-applied:

Refer to instructions for use of **STRELIUS II** under **APPLICATION PROCEDURES** section on this label. For minimum-tillage or no-tillage systems only, **StreliaS II** + AAtrex may be applied up to 45 days prior to planting in IA, IL, eastern KS, MO, NE, and SD. Use only split applications for treatments made 30 - 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 1.5 pts./A of **StreliaS II** + 1.7 - 2.0 lbs./A of AAtrex Nine-0* on *medium soils* with 1.5% organic matter or greater. Apply 1.5 pts./A of **StreliaS II** + 1.7 - 2.0 lbs./A of AAtrex Nine-0 on *fine soils* with less than 1.5% organic matter, or apply 1.67 pts./A of **StreliaS II** + 2.0 - 2.2 lbs./A of AAtrex Nine-0 on *fine soils* with 1.5% organic matter or greater. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation after application is recommended to move **StreliaS II** + AAtrex into the soil.

Precautions: To avoid crop injury,

1. Do not use on coarse soils.
2. Do not use on medium soils with less than 1.5% organic matter.

Preplant Incorporated or Preemergence:

Refer to instructions for use of **STRELIUS II** under **APPLICATION PROCEDURES** on this label. On *medium soils* with 1.5 organic matter or greater, apply 1.0 pt./A of **StreliaS II** + 1.3 lbs./A of AAtrex Nine-0*. On *fine soils* with less than 1.5% organic matter, apply 1.0 pt./A of **StreliaS II** + 1.3 lbs./A of AAtrex Nine-0; on *fine soils* with 1.5% organic matter or greater, apply 1.2 - 1.33 pts./A of **StreliaS II** + 1.6 - 1.8 lbs./A of AAtrex Nine-0.

* When using AAtrex 4L, use equivalent rates. One lb. of AAtrex Nine-0 equals 1.8 pts. of AAtrex 4L.

Precautions: To avoid crop injury,

1. Do not use on coarse soils.
2. Do not use on medium soils with less than 1.5% organic matter.
3. Do not use in NM, OK, or TX, except in northeast OK and the TX Gulf Coast and Blacklands areas.
4. Do not apply preplant incorporated in AZ or the Imperial Valley of CA.

TANK MIXTURE OF STRELIUS II OR STRELIUS II+ AATREX, WITH GRAMOXONE BRANDS, LANDMASTER BW, TOUCHDOWN BRANDS, OR ROUNDUP BRANDS FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where sorghum (seed treated with Concep) is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone brands, Landmaster BW, Touchdown, or Roundup may be tank mixed with **StreliaS II** or **StreliaS II** + AAtrex. See **Comment No. 7** following **Table 2**. The **StreliaS II** or **StreliaS II** + AAtrex portion of the tank mixture provides preemergence control of the weeds listed on this label under the respective sections.

Refer to the label of each product used in combination and observe the planting details, restrictions, and all other precautions and limitations.

Application:

Apply before, during, or after planting, but before sorghum emerges. Add Gramoxone brands, Landmaster BW, Touchdown brands, or Roundup brands and apply as directed on the product labels.

Gramoxone Brands:

Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

Landmaster BW:

27 - 54 ozs./A depending on weed species and size. See the Landmaster BW label for weeds controlled, rates for specific weeds, and other information concerning use.

Touchdown Brands or Roundup Brands:

See the Touchdown brand or Roundup brand labels for weeds controlled, rates, and other use directions.



SWEET SORGHUM (SEED TREATED WITH CONCEP)

Apply **StreliuS II** preplant surface, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Apply **StreliuS II** only when the sweet sorghum seed has been properly treated with Concep seed treatment. Preplant or preemergence applications of **StreliuS II** to sweet sorghum not treated with Concep seed treatment will result in crop death.

Soil-applied:

Apply **StreliuS II** up to 45 days before planting. Use only split applications for treatments made 30 - 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation after application is recommended to move **StreliuS II** into the soil.

StreliuS II Rates for Soil Applications to Sweet Sorghum

Soil Type	30 - 45 Days Prior to Planting ¹	<30 Days Prior to Planting	At Planting ²
COARSE	Not Recommended	1.33 pts./A	1.0 - 1.33 pts./A
MEDIUM	1.5 pts./A	1.5 pts./A	1.33 - 1.5 pts./A
FINE	1.67 pts./A	1.67 pts./A	1.33 - 1.67 pts./A

¹ Use only as a split application with 2/3 of the broadcast rate applied initially and the remaining 1/3 applied at planting.

² Preplant incorporated or preemergence.

Post-applied:

StreliuS II may be applied postemergence to sweet sorghum for residual control of grasses and small-seeded broadleaf weeds. Postemergence application to sweet sorghum may be made to crop up to 5 inches in height. **StreliuS II** will not control emerged weeds. Therefore, emerged weeds must be controlled by cultural or other chemical methods. When applied alone, **StreliuS II** will be safe to emerged sweet sorghum. Use of adjuvants is prohibited on sweet sorghum.

StreliuS II Rates for Postemergence Applications to Sweet Sorghum

Soil Type	Postemergence Rate
COARSE	1.0 - 1.33 pts./A
MEDIUM	1.33 pts./A
FINE	1.33 pts./A

Precautions:

1. If sweet sorghum seed is not properly treated with Concep seed treatment, soil applications of **StreliuS II** prior to sorghum emergence will severely injure the crop.
2. Under high soil moisture conditions prior to sweet sorghum emergence, injury may occur following soil applications of **StreliuS II**. The crop will normally outgrow this effect.
3. Do not use **StreliuS II** on sorghum grown under dry mulch tillage, or injury may occur.

Restrictions:

1. Only one application per season is allowed. **StreliuS II** may be applied either as a soil-applied treatment or a postemergence treatment, but not both.
2. Do not apply **StreliuS II** postemergence within 90 days of harvest.

SOYBEANS STRELIUS II ALONE

Apply **StreliuS II** preplant surface-applied, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Follow instructions for use of **StreliuS II** alone under **APPLICATION PROCEDURES** section of this label.

Fall Application for Spring Weed Control:

1. Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
2. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
3. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 - 2.0 pts./A of **StreliuS II** on *medium-textured* and 2.0 pts./A of **StreliuS II** on *fine-textured* soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2 - 3 inches. Minimize furrow and ridge formation in the tillage operations.

Restrictions: If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for soybeans of 2.5 pts./A depending on soil texture, or illegal residues may result.

Fall Application for Italian Ryegrass Control:

StreliuS II may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply **StreliuS II** at 1.33 - 1.67 pts./A in the fall (September 1 - December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower **StreliuS II** rate for *coarse-textured* soils and the higher rate for *fine-textured* soils. A tillage operation may precede the application. Do not incorporate to a depth greater than 2 - 3 inches if tillage follows the application of **StreliuS II**. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone brands can be tank mixed with **StreliuS II** for control of emerged ryegrass. Refer to the Gramoxone brands label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with **StreliuS II** for control of improved control of other weeds present at the time of application.

Restrictions:

1. Do not apply **StreliuS II** to frozen ground.
2. If a spring application is made, the combined total amount of **StreliuS II** applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for soybeans (2.5 pts./A, depending on soil texture), or illegal residues may result.

Preplant Surface - Spring Application:

Use on *medium* and *fine* soils with minimum-tillage or no-tillage systems in CO, CT, DE, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, MT, ND, NE, NH, NY, OH, PA, RI, SD, TN, VA, VT, WI, WV, and WY. Apply 2/3 the listed rate of **StreliuS II** (1.67 pts./A on *medium* soils and 2.0 pts./A on *fine* soils) as a split treatment 30 - 45 days prior to planting and the remainder at planting. Applications made less than 30 days before planting may be as either a split or single treatment. Apply 1.33 pts./A of **StreliuS II** on *coarse* soils not more than 2 weeks prior to planting.

Restrictions:

1. On soybeans, use up to 2.5 pts./A of **StreliuS II** preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%.
2. The total **StreliuS II** rate applied to soybeans during any one crop should not exceed 2.5 pts./A.
3. Do not graze or feed treated soybean forage, hay, or straw to livestock 30 days following treatment, or illegal residues may result.



Preplant Incorporated or Preemergence:

On *coarse soils*, apply 1.0 - 1.33 pts./A of **Strelius II** if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts./A of **Strelius II**. On *fine soils*, apply 1.33 - 1.67 pts./A of **Strelius II** if organic matter content is less than 3%, or 1.67 - 2.0 pts./A if organic matter content is 3% or greater.

Restrictions:

1. On soybeans, use up to 2.5 pts./A of **Strelius II** preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%.
2. The total **Strelius II** rate applied to soybeans during any one crop should not exceed 2.5 pts./A.
3. Do not graze or feed treated soybean forage, hay, or straw to livestock 30 days following treatment, or illegal residues may result.

Postemergence:

Apply 1.0 - 1.33 pts./A as a postemergence treatment to soybeans from emergence up through the third trifoliate leaf stage. **Strelius II** will not control emerged weeds so it must be applied to a weed-free soil surface or in a tank mixture with products that provide postemergence control of weeds present at the time of application.

Strelius II can also be applied as part of a sequential soybean weed control program. If **Strelius II** was applied as a preplant surface, preplant incorporated, or a preemergence treatment, a second treatment of **Strelius II** can be applied postemergence provided that the total **Strelius II** rate during any one crop does not exceed 2.5 pts./A.

Restrictions: To avoid possible illegal residues when **Strelius II** is applied postemergence to soybeans:

1. Do not apply more than 1.33 pts./A postemergence.
2. The total **Strelius II** rate applied preplant, preemergence or postemergence to soybeans during any one crop should not exceed 2.5 pts./A (2.4 lbs. a.i./A of *S-metolachlor*).
3. Make postemergence application at least 90 days before harvest.
4. Do not graze or feed treated forage or hay from soybeans to livestock following a postemergence application of **Strelius II**.

SOYBEANS

STRELIUS II COMBINATIONS

Water or fluid fertilizer may be used as carrier for **Strelius II** in combination with Sencor, Lorox, Canopy, Pursuit, Scepter, Sonalan, or Command.

Restrictions: For all of the following combinations, on soybeans use up to 2.5 pts./A **Strelius II** preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%. The total **Strelius II** rate applied to soybeans during any one crop year should not exceed 2.5 pts./A.

TANK MIXTURE WITH SENCOR

In addition to those weeds controlled by **Strelius II** alone, **Strelius II** + Sencor, when applied as directed, also controls the following broadleaf weeds: cocklebur*, hairy nightshade, hemp sesbania, jimsonweed*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard.

* Partially controlled.

Apply **Strelius II** and Sencor preplant incorporated or preemergence, using the appropriate rates from **Table 11. Preplant Incorporated or Preemergence:** Follow instructions for use of **Strelius II** alone under **APPLICATION PROCEDURES.**

Sequential:

Apply **Strelius II** alone **Preplant Incorporated**, as specified in **Table 11** for this tank mixture. Follow with a preemergence application of Sencor during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

Refer to the Sencor label for planting details and soybean variety restrictions.

Table 11: Strelius II + Sencor - Soybeans

Soil Texture**	Broadcast Rates Per Acre	
	0.5% to Less Than 3% Organic Matter	3% Organic Matter or Greater
	Strelius II + Sencor*	Strelius II + Sencor*
COARSE Loamy sand (over 2% organic matter), sandy loam	0.8 - 1.0 pt. + 0.33 lb.	1.0 pt. + 0.5 lb.
MEDIUM	1.0 - 1.33 pts. + 0.5 lb.	1.33 pts. + 0.67 lb.
FINE	1.33 pts. + 0.67 lb.	1.33 - 1.67 pts. + 0.67 lb.
Mississippi Delta only Silty clay, clay	1.33 pts. + 1.0 lb.	1.33 - 1.67 pts. + 1.0 lb.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE	

* When using Sencor 4, multiply lbs. of DF by 1.5 to get pts./A.

** On all sand and on loamy sand with less than 2% organic matter, do not use this tank mixture preemergence, or the sequential treatment. Do not use the tank mixture preplant incorporated on any sand, loamy sand, or sandy loam, or crop injury may occur.

*** Use 0.5 lb./A if applied preplant incorporated.

Precautions:

1. Do not use the tank mix or sequential application on soil with less than 0.5% organic matter or on alkaline soil with a pH over 7.4, or crop injury may occur.
2. If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days.

Restrictions: Follow most restrictive limitations and precautions on the **SOYBEANS - STRELIUS II ALONE** section of the **Strelius II** label and the soybean directions on the Sencor label.

TANK MIXTURE WITH LOROX

In addition to those weeds controlled by **Strelius II** alone, **Strelius II** + Lorox, applied preemergence, also controls the following broadleaf weeds: cocklebur*, jimsonweed*, lambsquarters, morningglory*, prickly sida, ragweed, smartweed, velvetleaf*, Venice mallow, and wild mustard.

* Partially controlled.



Preemergence: Apply during planting (behind planter) or after planting, but before weeds or soybeans emerge. Refer to the Lorox label for planting details. Apply the appropriate rates from **Table 12**.

Precaution: Do not use on soil with less than 0.5% organic matter, or crop injury may occur.

Table 12: Strelius II + Lorox - Soybeans

Soil Texture*	Broadcast Rates Per Acre	
	0.5% to Less Than 3% Organic Matter	3% Organic Matter or Greater
	Strelius II + Lorox DF	Strelius II + Lorox DF
COARSE**	0.8 pt. + 1.0 lb.	1.0 pt. + 1.0 - 1.5 lbs.
MEDIUM	1.0 pt. + 1.0 - 1.5 lbs.	1.33 pts. + 1.5 - 2.0 lbs.
FINE	1.33 pts. + 2.0 lbs.	1.33 - 1.67 pts. + 2.5 - 3.0 lbs.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE	

* Do not use on sand, gravelly soils, or exposed subsoils.

** Do not use on loamy sand, except in the northeastern U.S. on loamy sand with over 1% organic matter.

TANK MIXTURE WITH TREFLAN

Strelius II + Treflan tank mix applied preplant incorporated controls those weeds listed under **STRELIUS II APPLIED ALONE** and those weeds listed for Treflan Alone on the Treflan label. **Strelius II + Treflan** may be applied by ground or aerial equipment and incorporated up to 14 days before planting. Follow the recommended procedures on the Treflan and **Strelius II** labels, using equipment that provides uniform 2-inch incorporation.

Apply **Strelius II + Treflan** tank mix using the appropriate rate from the **SOYBEANS - STRELIUS II ALONE** section of this label and the Treflan Alone section of the Treflan label for the specific soil texture/organic matter classification and weed species expected.

To control DNA-resistant goosegrass* and other species on the respective labels where the soil organic matter is 3% or less, apply the rate in **Table 13**.

Table 13: Strelius II + Treflan - Organic Matter Content Less Than 3%

Soil Texture	Broadcast Rates Per Acre		
	Strelius II Organic Matter Less Than 3%	Treflan E.C.** Organic Matter	
		Less Than 2%	2 - 3%
COARSE*	0.8 - 1.0 pt.	1.0 pt.	1.5 pts.
MEDIUM	1.0 pt.	1.5 pts.	1.5 pts.
FINE	1.33 pts.	2.0 pts.	2.0 pts.

* Where a range of rates is given for **Strelius II**, use the minimum rate where DNA-resistant goosegrass is the predominant species.

** When Treflan 5 is used, use comparable rates. Multiply pts. of Treflan E.C. by 0.8 for Treflan 5.

Restrictions: Follow the most restrictive limitations and precautions on the **SOYBEANS - STRELIUS II ALONE** section of the **Strelius II** label and the soybean directions on the Treflan labels.

TANK MIXTURE WITH SCEPTER

This tank mixture controls all weeds controlled by **Strelius II** alone and by Scepter alone. Refer to the **STRELIUS II APPLIED ALONE** section for weeds controlled by **Strelius II** and to the Scepter label for weeds controlled by Scepter. Refer to the Scepter label for geographical locations where this tank mixture may be applied.

Apply **Strelius II + Scepter** preplant incorporated or preemergence, using rates in **Table 14**. Follow use directions under **Application Instructions** on the Scepter label. For preplant incorporated applications, apply and incorporate within 30 days before planting. Observe all other precautions and limitations on the Scepter labels.

Table 14: Strelius II + Scepter - Soybeans

Soil Texture	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	Strelius II	Scepter	Strelius II	Scepter
COARSE	0.8 pt.	0.67 pt.	1.0 pt.	0.67 pt.
MEDIUM	1.0 pt.	0.67 pt.	1.33 pts.	0.67 pt.
FINE	1.33 pts.	0.67 pt.	1.33 - 1.67* pts.	0.67 pt.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE			

* Use the higher rate of **Strelius II** if heavy weed infestations are expected.

Restrictions: Follow the most restrictive limitations and precautions on the **SOYBEANS - STRELIUS II ALONE** section of the **Strelius II** label and the soybean directions on the Scepter label.

TANK MIXTURE WITH CANOPY

This tank mixture controls all weeds controlled by **Strelius II** alone and by Canopy alone. Refer to the **STRELIUS II APPLIED ALONE** section for weeds controlled by **Strelius II** and to the Canopy label for weeds controlled by Canopy.

Apply preplant incorporated or preemergence, using the appropriate rates from **Table 15**. **Preplant Incorporated:** Apply within 2 weeks of planting. Uniformly incorporate into the top 1 - 2 inches of soil before planting soybeans. **Preemergence:** Apply after planting, but before soybeans emerge.

Restrictions: Follow the most restrictive limitations and precautions on the **SOYBEANS - STRELIUS II ALONE** section of the **Strelius II** label and the soybean directions on the Canopy label, including varietal restrictions.



Table 15: Strelius II + Canopy - Soybeans

Soil Texture	Broadcast Rates Per Acre		
	Less Than 3% Organic Matter		3% or More Organic Matter
	Strelius II	Strelius II	Canopy
COARSE	0.8 pt.	1.0 pt.	*
MEDIUM	1.0 pt.	1.33 pts.	*
FINE	1.33 pts.	1.33 - 1.67 pts.	*

* Refer to the Canopy label for appropriate rate according to geographical location, soil and organic matter classification, and pH limitations.

Precaution:

1. Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Canopy label.

TANK MIXTURE WITH COMMAND*

This tank mixture controls all weeds controlled by **Strelius II** alone and by Command alone. Refer to the **STRELIUS II APPLIED ALONE** section for weeds controlled by **Strelius II** and to the Command label for weeds controlled by Command.

Apply **Strelius II** + Command preplant incorporated, using rates in **Table 16**. Follow all Command application instructions as to incorporation interval, geographical location, equipment operation, soil moisture conditions, etc.

* **Restrictions:** Follow the most restrictive limitations and precautions on the **SOYBEANS - STRELIUS II ALONE** section of the **Strelius II** label and the soybean directions on the Command label, including rotational restrictions.

Table 16: Strelius II + Command - Soybeans

Soil Texture	Broadcast Rates Per Acre	
	Strelius II	
	0.5 - 3% Organic Matter	Greater Than 3% Organic Matter
COARSE	0.8 pt.	1.0 pt.
MEDIUM	1.0 pt.	1.33 pts.
FINE	1.33 pts.	1.33 - 1.67 pts.

TANK MIXTURE WITH SONALAN

This tank mixture controls all weeds controlled by **Strelius II** alone and by Sonalan alone. Refer to the **STRELIUS II APPLIED ALONE** section for weeds controlled by **Strelius II** and to the Sonalan label for weeds controlled by Sonalan.

Apply **Strelius II** and Sonalan preplant incorporated, using the appropriate rates from **Table 17**.

Preplant Incorporated: Follow recommended soil preparation procedures for Sonalan.

Sequential: Apply Sonalan alone preplant incorporated as specified on the Sonalan label. Follow with a preemergence application of **Strelius II** during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

Table 17: Strelius II + Sonalan - Soybeans

Soil Texture	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	Strelius II	Sonalan	Strelius II	Sonalan
COARSE	1.0 - 1.33 pts.	1.25 - 2.0 pts.	1.33 pts.	1.25 - 2.0 pts.
MEDIUM*	1.33 - 1.67 pts.	1.75 - 2.5 pts.	1.33 - 1.67 pts.	1.75 - 2.5 pts.
FINE*	1.33 - 1.67 pts.	2.25 - 3.0 pts.	1.67 - 2.0 pts.	2.25 - 3.0 pts.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE			

* For eastern black nightshade on these soils, apply Sonalan at 3.0 pts./A on *medium* and 3.5 pts./A on *fine-textured soils*, and follow with 2 incorporation passes.

Restrictions: Follow the most restrictive limitations and precautions on the **SOYBEANS - STRELIUS II ALONE** section of the **Strelius II** label and the soybean directions on the Sonalan label.

TANK MIXTURE WITH PURSUIT

This tank mixture controls all weeds controlled by **Strelius II** alone and by Pursuit alone. Refer to the **STRELIUS II APPLIED ALONE** section for weeds controlled by **Strelius II** and to the Pursuit label for weeds controlled by Pursuit. Refer to the Pursuit label for geographical locations where this tank mixture may be applied.

Apply **Strelius II** + Pursuit early preplant, preplant incorporated, or preemergence after planting, using rates in **Table 18**. Application can be made in water or liquid fertilizer. Follow all use directions under Soil Applications on the Pursuit label. For early preplant and preplant incorporated applications, apply within 30 days before planting.

Restrictions: Follow the most restrictive limitations and precautions on the **SOYBEANS - STRELIUS II ALONE** section of the **Strelius II** label and the soybean directions on the Pursuit label, including rotational restrictions.

Table 18: Strelius II + Pursuit - Soybeans

Soil Texture	Broadcast Rates Per Acre		
	Less Than 3% Organic Matter	3% or More Organic Matter	Less Than 3% - 3% or More Organic Matter
	Strelius II	Strelius II	Pursuit
COARSE	0.8 pt.	1.0 pt.	0.25 pt.
MEDIUM	1.0 pt.	1.33 pts.	0.25 pt.
FINE	1.33 pts.	1.33 - 1.67 pts.	0.25 pt.

Sequential:

Apply **Strelius II** early preplant, preplant incorporated, or preemergence after planting at 0.8 pt./A on *coarse soils* and 1.0 pt./A on *medium- and fine-textured soils*. Follow with a sequential postemergence application of Pursuit to control emerged weeds according to the Pursuit label. **Strelius II** will improve the consistency and level of control from Pursuit on most grass species. Refer to the Pursuit postemergence label for a listing of weeds controlled, application rate, and growth stage limitations.



TANK MIXTURE WITH SENCOR, SCEPTER, LOROX, CANOPY, OR PURSUIT, PLUS GRAMOXONE BRANDS, TOUCHDOWN BRANDS, OR ROUNDUP BRANDS FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where soybeans are planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone brands, Touchdown brands, or Roundup brands may be added to a tank mix of either **Strelius II** + Sencor, **Strelius II** + Scepter, **Strelius II** + Lorox, **Strelius II** + Canopy, or **Strelius II** + Pursuit. When used as directed, the Gramoxone brands portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Touchdown or Roundup combinations will control emerged annual and perennial weeds when applied as directed on the Touchdown or Roundup label. The **Strelius II** + Sencor, Scepter, Lorox, Canopy, or Pursuit portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for **Strelius II** + Sencor, **Strelius II** + Scepter, **Strelius II** + Lorox, **Strelius II** + Canopy, and **Strelius II** + Pursuit, respectively.

Refer to the label of each product used in combination and observe the planting details, soybean variety restrictions, information regarding application to soybeans, geographical restrictions, and all other precautions and limitations.

Application:

Apply before, during, or after planting, but before the soybeans emerge. Add Gramoxone brands, Touchdown brands, or Roundup brands and apply as directed on the product labels.

Gramoxone Brands:

Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

Restrictions: Do not apply combinations containing Gramoxone brands in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

Touchdown or Roundup:

See the Touchdown brand or Roundup brand label for weeds controlled, rates, and other use directions.

Apply in 20 - 60 gals. of water or fluid fertilizer per acre with ground equipment.

Strelius II + Sencor + Gramoxone Brands, Touchdown Brands, or Roundup Brands

On loamy sand with over 2% organic matter, apply 1.0 pt./A of **Strelius II** + 0.33 - 0.5 lb./A of Sencor. On *medium soils*, apply 1.33 pts./A of **Strelius II** + 0.5 - 0.67 lb./A of Sencor. On *fine soils*, apply 1.33 - 1.67 pts./A of **Strelius II** + 0.67 lb./A of Sencor.

* When using Sencor 4 or Lexone® 4L, multiply lbs. of DF by 1.5 to get pts./A.

Precautions: To avoid crop injury,

1. Do not use this tank mixture on soil with less than 0.5% organic matter, on alkaline soil with a pH over 7.4, or on all sand and on loamy sand with less than 2% organic matter.
2. If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.

Strelius II + Scepter + Gramoxone Brands, Touchdown Brands, or Roundup Brands

On *coarse soils*, apply 1.0 pt./A of **Strelius II** + 0.67 pt./A of Scepter. On *medium soils*, apply 1.33 pts./A of **Strelius II** + 0.67 pt./A of Scepter. On *fine soils*, apply 1.67 pts./A of **Strelius II** + 0.67 pt./A of Scepter.

Restrictions: (1) Do not apply within 90 days of harvest, and (2) Do not graze or feed treated soybean forage, hay, or straw to livestock, or illegal residues may result.

Strelius II + Lorox + Gramoxone Brands, Touchdown Brands, or Roundup Brands

On *coarse soils**, apply 1.0 pt./A of **Strelius II** + 1.0 - 1.5 lbs./A of Lorox DF. On *medium soils*, apply 1.33 pts./A of **Strelius II** + 1.0 - 2.0 lbs./A of Lorox DF. On *fine soils*, apply 1.33 - 1.67 pts./A of **Strelius II** + 2.0 - 3.0 lbs./A of Lorox DF.

* Do not use on loamy sand, except in the northeastern U.S. on loamy sand with over 1% organic matter, or injury may occur. Do not use on sand, gravelly soils, or exposed subsoils, or injury may occur.

Precaution: Do not use on soil with less than 0.5% organic matter, or crop injury may occur.

Strelius II + Canopy + Gramoxone Brands, Touchdown Brands, or Roundup Brands

Use only where soils have 0.5 - 5% organic matter. On *coarse soils* (except sand), apply 1.0 pt./A of **Strelius II**, on *medium soils*, apply 1.33 pts./A of **Strelius II**, and on *fine soils*, apply 1.33 - 1.67 pts./A of **Strelius II**. Refer to the Canopy label for appropriate rate, according to geographical location, soil and organic matter classification, pH limitations, and all other use directions.

Precaution: Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Canopy label.

Strelius II + Pursuit + Gramoxone Brands, Touchdown Brands, or Roundup Brands

On *coarse soils*, apply 1.0 pt./A of **Strelius II** + 0.25 pt./A of Pursuit. On *medium soils*, apply 1.33 pts./A of **Strelius II** + 0.25 pt./A of Pursuit. On *fine soils*, apply 1.67 pts./A of **Strelius II** + 0.25 pt./A Pursuit.

POSTEMERGENCE USE ON SOYBEANS - STRELIUS II TANK MIXTURES

Tank Mixture with Glyphosate Products

(e.g., Touchdown Brands or Roundup Brands)

Strelius II at 1.0 - 1.33 pts./A may be tank mixed with glyphosate products at labeled rates and applied from emergence up through the third trifoliate leaf stage of glyphosate-tolerant soybeans. **Strelius II** alone will not control emerged weeds. Use this treatment only on soybeans designated for use with glyphosate (e.g., glyphosate-tolerant soybeans). The glyphosate product must be registered for postemergence use in glyphosate-tolerant soybeans.

Tank Mixture with Pursuit

Strelius II at 1.0 - 1.33 pts./A may be tank mixed with Pursuit at labeled rates and applied from emergence up through the third trifoliate leaf stage of soybeans. **Strelius II** alone will not control emerged weeds.

Tank Mixture with Liberty Herbicide or Ignite 280 SL

Strelius II at 1.0 - 1.33 pts./A may be tank mixed with Liberty Herbicide or Ignite 280 SL Herbicide at labeled rates and applied from emergence up through the third trifoliate leaf stage of soybeans. **Strelius II** alone will not control emerged weeds. Use this treatment only on soybeans designated for use with glufosinate (e.g., LibertyLink).

Restrictions: Follow the tank mix product label for adjuvant recommendations. The use of COC or UAN with **Strelius II** may result in temporary crop injury. To avoid possible illegal residues when **Strelius II** is applied post-emergence to soybeans:

1. Do not apply more than 1.33 pts./A postemergence.
2. Make postemergence application at least 90 days before harvest.
3. Do not graze or feed treated forage or hay from soybeans to livestock following a postemergence application of **Strelius II**.



SUGAR BEETS

STRELIUS II ALONE

Postemergence Applications

Strelius II may be applied postemergence to sugar beets after the sugar beets have reached the first true-leaf stage. However, because **Strelius II** is primarily a soil-active herbicide, it must be applied prior to weed emergence in order to provide consistent control of listed weeds. As such, weeds that are emerged with or before the crop, or that are present at the time **Strelius II** is applied, must be controlled with another appropriately labeled herbicide. Apply **Strelius II** at 1 pt./A on *coarse soils*, 1.33 pts./A on *medium soils*, and 1.67 pts./A on *fine soils*. More than one postemergence application may be applied, but the total should not exceed 2.6 pts./A. Weeds present at the time of application will not be controlled.

Restrictions: To avoid possible illegal residues:

1. Do not apply more than 2.6 pts./A postemergence.
2. Do not harvest within 60 days after the last application.

Precaution: In *coarse soils*, **Strelius II** applied before emergence of sugar beets (i.e., other than postemergence) may cause injury.

SUGAR BEETS

STRELIUS II TANK MIX COMBINATIONS

Strelius II may be tank mixed with Assure® II, Betamix®, Poast®, Progress®, Stinger®, or Upbeet® and applied to sugar beets. Tank mixtures of these products with **Strelius II** will increase the risk of crop injury over that of either product applied alone, as the **Strelius II** formulation has some adjuvant properties. The addition of a spray adjuvant such as crop oil concentrates (COC's) or methylated seed oils (MSO's) can further increase the risk of crop injury. Injury risk can be reduced by using the lowest effective rate of the tank mix partner(s) and/or adjuvant and by avoiding applications under adverse growing conditions or high soil or air humidity. Refer to the individual product labels and follow all use restrictions and limitations.

SUGARCANE*

STRELIUS II ALONE

*Not for Use in California

Strelius II can be applied as a preplant, preemergence, or postemergence treatment for weed control in sugarcane. **Strelius II** can also be used in a treatment program that includes a preplant/preemergence application followed by a postemergence/post-directed application.

Preplant and Preemergence Applications

Apply **Strelius II** at a rate of 0.89 – 1.22 qt/A prior to planting, preemergence after new plantings, or after harvest, but prior to re-emergence of ratoon-cane. Apply the higher rate on heavier soils and soils with higher organic matter content. For a list of weeds controlled preemergence, refer to Table 1.

If weeds are emerged at the time of application, add a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v (1 gallon/100 gallon) **or** a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v (1 quart/100 gallon) to the spray solution. In addition to COC or NIS, a spray grade UAN (e.g. 28-0-0) at a rate of 2.5% v/v (2.5 gal/100 gal spray solution) **or** ammonium sulfate (AMS) at a rate of 8.5 lb/100 gallons of spray solution can be added to the spray solution.

Postemergence Applications

Apply **Strelius II** at a rate of 0.49 - 0.98 qt/A postemergence before the sugarcane reaches 60 inches in height. Refer to Table 1 for a list of weeds controlled.

Postemergence applications may be made as broadcast post-over-the-top or as a post-directed spray to soil between the rows and the base of the sugarcane. For best postemergence weed control, apply **Strelius II** to actively growing weeds.

If a preemergence application was made earlier in the season (not to exceed 1.22 qt/A), only 0.49 qt/A may be applied postemergence. The total amount of **Strelius II** applied (preemergence + postemergence) cannot exceed 1.71 qt/A/year.

If weeds are emerged at the time of application, add a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v (1 gallon/100 gallon) **or** a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v (1 quart/100 gallon) to the spray solution. In addition to COC or NIS, a spray grade UAN (e.g. 28-0-0) at a rate of 2.5% v/v (2.5 gal/100 gal spray solution) **or** ammonium sulfate (AMS) at a rate of 8.5 lb/100 gallons of spray solution can be added to the spray solution.

Strelius II Tank Mix Combinations

Strelius II may be tank mixed with other registered sugarcane herbicides, insecticides and fungicides. The tank-mix partners are to be applied by the same methods and at the same timings as **Strelius II** unless otherwise specified in the tank-mix product label. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Perform a compatibility test prior to spraying the tank-mix application.

Restrictions: To avoid possible illegal residues:

1. Do not apply more than 1.71 qt/A of **Strelius II** per year.
2. Do not apply more than 1.22 qt/A of **Strelius II** as a preplant or preemergence application per year.
3. Do not apply more than 0.98 qt/A of **Strelius II** as a postemergence application per year.
4. Do not apply more than 0.49 qt/A of **Strelius II** as a postemergence application if a preplant or preemergence application was made.
5. Do not make more than two applications per year.
6. Do not make applications less than 14 days apart.
7. Do not apply **Strelius II** to sugarcane greater than 60 inches in height.
8. Do not make a postemergence application within 100 days of harvest.
9. Do not exceed the total combined maximum annual sugarcane rates for S-metolachlor containing products (3.34 lb S-metolachlor ai/A).

SUNFLOWERS

STRELIUS II ALONE

Preplant Incorporated or Preemergence

Within the rate ranges given below, use the higher rate of **Strelius II** if heavy weed infestations are expected. On *coarse soils* with organic matter less than 3%, apply 1.0 - 1.33 pts./A of **Strelius II**; apply 1.33 pts./A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts./A of **Strelius II**. On *fine soils* with organic matter of less than 3%, apply 1.33 - 1.67 pts./A of **Strelius II**; apply 1.67 - 2.0 pts./A if organic matter is 3% or greater.

Restrictions:

1. Do not allow livestock to graze or feed in treated area.
2. Do not exceed the maximum label rates for sunflowers for the soil type.



TOMATOES

STRELIUS II ALONE

Transplanted

Strelus II may be applied preplant incorporated or preplant before transplanting. If the latter method is used, keep soil disturbance to a minimum during the transplanting operation. Application may also be post directed to transplants after the first settling rain or irrigation. When an application is made post-directed, apply in a minimum of 20 gallons of water per acre and minimize contact with tomato plants. **Strelus II** will not control emerged weeds. In bedded transplanted tomatoes, apply **Strelus II** preplant non-incorporated to the top of the pressed bed as the last step prior to laying plastic. **Strelus II** may also be used to treat row-middles in bedded tomatoes, as long as the total amount of **Strelus II** does not exceed the maximum allowed per crop.

Seeded

Strelus II may be applied post-directed to direct-seeded tomatoes: Tomato plants must be at least 4 inches tall at the time of application, and the product must be applied in a minimum of 20 gallons of water per acre. Minimize spray contact with tomato plants. **Strelus II** will not control emerged weeds.

Tomato Use Rates:

On *coarse soils*, apply 1.0 - 1.33 pts./A of **Strelus II** if organic matter is less than 3% or 1.33 pts./A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts./A of **Strelus II**. On *fine soils*, apply 1.33 - 1.67 pts./A of **Strelus II** if organic matter is less than 3% or 1.67 - 2.0 pts./A if organic matter is 3% or greater.

Precautions:

- Strelus II** may damage transplants that have been weakened by any cause. To prevent damage, plant only healthy transplants. Do not plant when wet, cool, or unfavorable growing conditions exist.
- In transplanted tomatoes, if **Strelus II** is applied preplant incorporated, incorporate to a depth less than the depth of transplanting, and use the lower end of the rate range for the given soil type, or damage may occur.
- For row-middle applications where tomatoes are grown on sandy soils and where high soil moisture conditions can exist (e.g., low binding and high evaporation conditions), as may be found in the States of Florida, Georgia, Maryland, and Virginia, there is potential for crop injury in the form of leaf epinasty. The risk of this type of injury can be reduced by: a) incorporating the **Strelus II** immediately following application, b) applying the **Strelus II** seven or more days before transplanting (but only after the beds have been formed), c) minimizing the application of **Strelus II** onto the plastic of the bed, or d) any combination of the above.

Restrictions:

- Do not apply to varieties or cultivars with unknown tolerance to **Strelus II**.
- 90-Day PHI - If the single application rate of **Strelus II** is greater than 1.33 pts./A (up to 2.0 pt./A), do not harvest tomatoes within 90 days of application.
- 30-Day PHI - If the application of **Strelus II** does not exceed 1.33 pts./A, do not harvest tomatoes within 30 days of application.
- When applying at 1.33 pts./A with a 30-day PHI, the following restrictions apply:
 - Do not exceed two applications per growing season.
 - The use of adjuvants is prohibited.
 - Applications may be made using ground equipment, in concentrated spray volumes.
 - Applications may be made as a foliar broadcast spray to the soil within 1 week of transplanting and again at blooming/fruiting to the row middles as a banded/directed application 38-77 days after the first treatment.

To avoid possible illegal residues:

- Do not exceed the maximum label rate for the soil texture per year.
- Apply only by ground application.

Tank Mix Partner Information

Product Name	Active Ingredient	EPA Registration Number
AAtrex 4L	atrazine	100-497
AAtrex Nine-O	atrazine	100-585
Accent	nicosulfuron	352-560
Assure II	quizalofop-p-ethyl	5481-646
Balance Pro	isoxaflutole	264-600
Banvel®	dicamba	66330-276
Basagran®	sodium bentazon	7969-45
Beacon	primisulfuron-methyl	100-705
Betamix	desmedipham	264-621
Bicep Magnum	atrazine + S-metolachlor	100-886
Bicep II Magnum	atrazine + S-metolachlor	100-817
Buctril	bromoxynil	264-437
Butyrac® 200	butyric acid, dimethylamine salt	42750-38
Canopy®	metribuzin + chlorimuron ethyl	352-444
Caparol® 4L	prometryn	100-620
Clarity	diglycolamine salt	7969-137
Command	clomazone	279-3158, 279-3150, 279-3151
Concep	fluxofenim	
Cotoran 4L	fluometuron	66222-181
Eptam® 7E	S-ethyl dipropylthiocarbamate	10163-283
Eradicane	S-ethyl dipropylthiocarbamate	10163-284
Gramoxone brands	paraquat	100-1652, 100-1431
Ignite 280 SL Herbicide	glufosinate	7969-448
Landmaster BW	glyphosate + 2,4-D	524-351
Lexone 4L	metribuzin	352-382

(continued)



Tank Mix Partner Information (cont.)

Product Name	Active Ingredient	EPA Registration Number
Liberty® Herbicide	glufosinate	7969-447
Lorox DF	linuron	61842-23
Marksman®	atrazine + dicamba	7969-136
Poast	sethoxydim	7969-176
Princep 4L	simazine	100-526
Princep Caliber 90	simazine	100-603
Progress	phenmedipham + desmedipham + ethofumesate	264-632
Prowl	pendimethalin	241-418
Pursuit®	imazethapyr	241-310
Roundup brands	glyphosate	524-445, 524-343, 524-539
Scepter® 70 DG	imazaquin	5481-610
Sencor 4	metribuzin	264-735
Sonalan	ethalfuralin	10163-356
Stinger	clopyralid	62719-73
Storm	sodium salt of bentazon + sodium salt of acifluorfen	70506-59
Touchdown brands	glyphosate	100-1169, 100-1157, 100-1212
Treflan	trifluralin	10163-360
Upbeet	triflusalufuron-methyl	279-9584

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: This product may be stored at temperatures down to -30°F.

Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

CONTAINER HANDLING

Nonrefillable Containers (≤ 5 gallons): Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Containers (> 5 gallons): Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure 2 more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Strelis™ is a trademark of Atticus, LLC

AAtrex®, AAtrex Nine-0®, Beacon®, Bicep Magnum®, Bicep II Magnum®, Caparol®, Concep®, Evik®, Gramoxone®, Princep®, Princep® Caliber 90®, Touchdown® are trademarks of a Syngenta Group Company

Accent®, Canopy®, Lexone®, and Viton® are trademarks of E. I. duPont de Nemours and Company, Inc.

Ag-Chem RoGator® trademark of Ag-Chem Equipment Company

Agsorb® trademark of Oil-Dri Corporation

Stinger® and Treflan® are trademarks of Dow AgroSciences

Balance® Pro, Betamix®, Buctril®, Butyrac®, Ignite®, Liberty®, Liberty® ATZ, LibertyLink®, Progress®, and Sencor® are trademarks of Bayer CropScience

Basagran®, Clarity®, Marksman®, Poast®, Prowl®, Pursuit®, and Storm® are trademarks of BASF Ag Products

Celatom MP-79® trademark of Eagle-Picher Industries, Inc.

Command® trademark of FMC Corporation

Compex® trademark of KALO Agricultural Chemicals, Inc.

Cotoran® trademark of Makhteshim Agan of North America, Inc.

Hi-Cycle™ trademark of John Deere Company

Landmaster®, Roundup®, and Roundup Ready® are trademarks of Monsanto Company

Tyler Patriot™ trademark of Tyler Ltd. Partnership

Unite® trademark of HACO, Inc.

Weedone® trademark of NuFarm, Inc.

Willmar Air Ride® trademark of Willmar Manufacturing

X-77® trademark of Loveland Industries, Inc.

20210623a

