

---

**SUPPLEMENTAL LABELING**

---

**Syngenta Crop Protection, LLC**

P. O. Box 18300

Greensboro, North Carolina 27419-8300

**SCP 1476A-S3 0922**

<b>BENZOVINDIFLUPYR</b>	<b>GROUP</b>	<b>7</b>	<b>FUNGICIDE</b>
<b>DIFENOCONAZOLE</b>	<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>

**Aprovia® Top Fungicide**

SOLATENOL® Technology\*

**This supplemental label expires on 09/23/2025 and must not be used or distributed after this date.**

## Active Ingredients:

Difenoconazole\*\* ..... 10.95%

Benzovindiflupyr\*\*\* ..... 7.30%

---

Other Ingredients: ..... 81.75%

---

Total: ..... 100.00%

\*Technology denotes the active ingredient, Benzovindiflupyr.

\*\*CAS No. 119446-68-3

\*\*\*CAS No. 1072957-71-1

Aprovia Top Fungicide is formulated as an emulsifiable concentrate containing 0.97 lb ai of difenoconazole active ingredient and 0.65 lb ai of benzovindiflupyr active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN.****WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

See First Aid Statement inside booklet and on container label.

EPA Reg. No. 100-1476

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Aprovia Top Fungicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**



## DIRECTIONS FOR USE

### Root Vegetable (Except Sugar Beet and Ginseng) Subgroup 1B

Crops (Including all cultivars, varieties, and/or hybrids of these) Not for use in California			
Beet, garden	Horseradish	Rutabaga	
Burdock, edible	Parsley, turnip-rooted	Salsify	
Carrot	Parsnip	Salsify, black	
Celeriac; chervil, turnip-rooted	Radish	Salsify, Spanish	
Chicory	Radish, oriental	Skirret; turnip.	
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Alternaria leaf spot ( <i>Alternaria dauci</i> ) Anthracnose ( <i>Colletotrichum spp.</i> ) Cercospora leaf spot ( <i>Cercospora carotae</i> ) Powdery Mildew ( <i>Erysiphe spp.</i> )	12.8*	Begin applications prior to disease onset when conditions are conducive for disease.	Apply by ground, air, or chemigation.  Apply on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.  The addition of a spreading/penetrating type adjuvant for example, a non-ionic based surfactant or crop oil concentrate or blend is advised.  If disease pressure is high, use the shortest interval.  See <b>Section 4.4.5</b> .  For chemigation, apply in 0.1 – 0.25 inches/A of water.
*12.8 fl oz product/A is equivalent to 0.065 lb ai benzovindiflupyr and 0.097 lb ai difenoconazole.			
<b>Resistance Management:</b>			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.2</b>.</li> </ul>			
<b>Precaution:</b>			
<ul style="list-style-type: none"> <li>Chemigation with excessive water may lead to a decrease in efficacy</li> </ul>			
USE RESTRICTIONS			
<ol style="list-style-type: none"> <li>Refer to <b>Section 6.1</b> for additional product use restrictions.</li> <li><b>Maximum Single Application Rate: DO NOT</b> exceed the maximum rate listed in the table.</li> <li><b>Minimum Application Interval:</b> 7 days</li> <li><b>Maximum Annual Rate:</b> 51.2 fl oz/A/year                             <ol style="list-style-type: none"> <li><b>DO NOT</b> exceed 0.26 lb ai/A/year of benzovindiflupyr-containing products.</li> <li><b>DO NOT</b> exceed 0.388 lb ai/A/year of difenoconazole-containing products.</li> </ol> </li> <li><b>DO NOT</b> exceed 4 application per year.</li> <li><b>DO NOT</b> apply by air in New York State.</li> </ol>			

7. **Pre-Harvest Interval (PHI):** 7 days

©2022 Syngenta

Aprovia® Top, SOLATENOL® and the Syngenta logo are trademarks of a Syngenta Group Company