

10/28/2021

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 5481-646

SUPPLEMENTAL USE DIRECTIONS**Assure II Herbicide**

EPA Reg. No. 5481- 646

KEEP OUT OF REACH OF CHILDREN**DANGER/ PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

Specific Crop Use Directions and Restrictions on Fruit, Pome (Crop Group 11-10), Fruit, Small Vine Climbing, except Fuzzy Kiwifruit (Subgroup 13-07F), Fruit, Stone (Crop Group 12-12), Carinata, Pennycress and Sunflower (subgroup 20B)

This label expires and must not be distributed or used after December 31st, 2023

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS SUPPLEMENTAL LABEL AND THE FULL PRODUCT LABEL MUST BE IN POSSESSION OF THE USER AT THE TIME OF APPLICATION

Read the complete label affixed to the container for Assure II before applying. Use of Assure II according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Assure II. This Supplemental Labeling contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Directions for Use

Specific Crop Use Directions and Restrictions

Apply ASSURE II to young, actively growing grasses according to the 'Weeds Controlled and Rate Selection' Table on the EPA registered label.

Fruit, Pome (Crop Group 11-10) (except New York state)

Assure® II is registered for control of annual and perennial grasses in Fruit, Pome Crop Group 11-10 [Apple; Azarole; Crabapple; Loquat; Mayhaw; Medlar; Pear; Pear, Asian; Quince; Quince, Chinese; Quince, Japanese; Tejocote; cultivars, varieties, and/or hybrids of these].

- ASSURE® II may be applied up to 12 fluid oz (0.0825 lb ai) per acre as a directed spray in a band extending out a minimum of 3 feet on each side of the tree row in 10-40 gallons of water/acre to control labelled grass weeds.
- Do not apply ASSURE® II within 14 days of harvest.
- Do not make more than 2 applications per acre per year.
- Application intervals must be at least 14 days apart to allow regrowth to occur.
- The maximum use rate is 24 fluid oz (0.165 lb ai) per acre per year.
- Do not allow the spray solution to contact the trees.
- Do not make aerial applications.
- Do not apply ASSURE II within 14 days of anticipated bloom of crop.

Fruit, Small Vine Climbing, except Fuzzy Kiwifruit (Subgroup 13-07F) (except New York state)

Assure® II is registered for control of annual and perennial grasses in Fruit, Small, Vine, Climbing, except Fuzzy Kiwifruit Crop Group 13-07F [Amur River Grape; Gooseberry; Grape; Kiwifruit, Hardy; Maypop; Schisandra Berry; cultivars, varieties, and/or hybrids of these].

- ASSURE® II may be applied at 5 fluid oz (0.034 lb ai) per acre as a directed spray in a band extending out a minimum of 3 feet on each side of the tree row in 10-40 gallons of water/acre to control labelled grass weeds.
- Do not apply ASSURE® II within 15 days of harvest.
- Do not make more than 2 applications per acre per year.
- Application intervals must be at least 14 days apart to allow regrowth to occur.
- The maximum use rate is 10 fluid oz (0.068 lb ai) per acre per year.
- Do not allow the spray solution to contact the vines.
- Do not make aerial applications.
 - Do not apply ASSURE II within 14 days of anticipated bloom of crop.

Fruit, Stone (Crop Group 12-12) (except New York state)

Assure® II is registered for control of annual and perennial grasses in Fruit, Stone Crop Group 12-12 [Apricot; Apricot, Japanese; Capulin; Cherry, Black; Cherry, Nanking; Cherry, Sweet; Cherry, Tart; Jujube, Chinese; Nectarine; Peach; Plum, American; Plum, Beach; Plum, Canada; Plum, Cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plum, Prune; Plumcot; Sloe; cultivars, varieties, and/or hybrids of these].

- ASSURE® II may be applied up to 12 fluid oz (0.0825 lb ai) per acre as a directed spray in a band extending out a minimum of 3 feet on each side of the tree row in 10-40 gallons of water/acre to control labelled grass weeds.
- Do not apply ASSURE® II within 14 days of harvest.
- Do not make more than 2 applications per acre per year.
- Application intervals must be at least 14 days apart to allow regrowth to occur.
- The maximum use rate is 24 fluid oz (0.165 lb ai) per acre per year.
- Do not allow the spray solution to contact the trees.
- Do not make aerial applications.
- Do not apply ASSURE II within 14 days of anticipated bloom of crop.

Carinata and Pennycress

ASSURE® II herbicide is registered for control of annual and perennial grasses in crops in Carinata and Pennycress.

- ASSURE® II may be applied from 4 - 12 fluid oz. (0.0275 – 0.0825 lb ai/A) in a minimum of 3 gallons of water per acre.
- Do not apply ASSURE® II within 60 days of harvest.
- The maximum use rate of ASSURE® II is 18 fluid oz (0.124 lb ai) per acre per year.
- Do not make more than 2 applications per acre per season.
- Application intervals must be greater than 7 days apart to allow regrowth to occur.
- Chemigation is prohibited.
- Do not apply ASSURE II within 14 days of anticipated bloom of crop.

Sunflower (Subgroup 20B)

ASSURE® II herbicide is registered for control of annual and perennial grasses in crops in Sunflower Subgroup 20B [Calendula; Caster Oil Plant; Chinese Tallowtree; Euphorbia; Evening Primrose; Jojoba; Niger Seed; Rose Hip; Safflower; Stokes Aster; Sunflower; Tallowwood; Tea Oil Plant; Vernonia; cultivars, varieties, and/or hybrids of these]

- ASSURE® II may be applied from 4 - 12 fluid oz. (0.0275 – 0.0825 lb ai/A) in a minimum of 3 gallons of water per acre.
- Do not apply ASSURE® II within 60 days of harvest.
- The maximum use rate of ASSURE® II is 18 fluid oz (0.124 lb ai) per acre per year.
- Do not make more than 2 applications per acre per season.
- Application intervals must be greater than 7 days apart to allow regrowth to occur.
- Chemigation is prohibited.
- Do not apply ASSURE II within 14 days of anticipated bloom of crop.

Assure is a trademark owned by AMVAC Chemical Corporation.

AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660