

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

GROUP

3

INSECTICIDE

Baythroid® XL

EPA Reg. No. 264-840

For use on Pome Fruit, Grape and Stone Fruit to control Spotted Wing Drosophila

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEW YORK

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

Pome Fruit Includes all members of Crop Group 11: Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental pear, and Quince			
PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb AI/Acre	Method
Spotted Wing Drosophila (adult)	2.4 – 2.8	0.019 – 0.022	Apply as a foliar spray, targeting adult growth stage.
Notes and Restrictions: Pre-Harvest Interval: 7 days Maximum BAYTHROID XL allowed per 14-day interval: 2.8 fluid ounces/Acre (0.022 lb AI/Acre) Maximum BAYTHROID XL allowed per crop season: 2.8 fluid ounces/Acre (0.022 lb AI/Acre)			

NYDEC SAL 11-02-11

Baythroid® XL EPA Reg. No. 264-840

For use on Pome Fruit, Grape and Stone Fruit to control Spotted Wing Drosophila

PRODUCT BULLETIN

Grape Includes: Table, Raisin, Muscadine, and Wine Grape			
PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb AI/Acre	Method
Spotted Wing Drosophila (adult)	2.4 – 3.2	0.019 – 0.025	Apply as a foliar spray, targeting adult growth stage. Reapply as necessary but no closer than a 14-day interval.
Notes and Restrictions: Pre-Harvest Interval: 3 days Maximum BAYTHROID XL allowed per 14-day interval: 3.2 fluid ounces/Acre (0.025 lb AI/Acre) Maximum BAYTHROID XL allowed per crop season: 12.8 fluid ounces/Acre (0.1 lb AI/Acre)			

Stone Fruit: Includes all members of Crop Group 12: Apricot, Cherry (sweet and tart), Nectarine, Peach, Plum, Plumcot, and Prune (fresh and dried)			
PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb AI/Acre	Method
Spotted Wing Drosophila (adult)	2.4 – 2.8	0.019 – 0.022	Apply as a foliar spray, targeting adult growth stage. Reapply as necessary but no closer than a 14-day interval.
Notes and Restrictions: Pre-Harvest Interval: 7 days Maximum BAYTHROID XL allowed per 14-day interval: 2.8 fluid ounces/Acre (0.022 lb AI/Acre) Maximum BAYTHROID XL allowed per crop season: 5.6 fluid ounces/Acre (0.044 lb AI/Acre)			

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937)*

NYDEC SAL 11-02-11