

Supplemental Labeling



FLONICAMID	GROUP	29	INSECTICIDE
------------	-------	----	-------------

SUPPLEMENTAL DIRECTIONS FOR THE USE OF
BELEAF® 50SG Insecticide
 (EPA Reg. No. 71512-10-279)

**FOR CONTROL OF LISTED PESTS ON
 GRAPES AND OTHER SMALL VINE CLIMBING FRUIT (EXCEPT FUZZY KIWIFRUIT) SUBGROUP 13-07F**

This Supplemental Label Expires on June 30, 2025 and must not be used or distributed after this date.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in possession of the user at the time of application. Read the label affixed to the container for BELEAF 50SG insecticide before applying. Use of BELEAF 50SG insecticide according to this labeling is subject to the use directions, restrictions and precautions imposed by the label affixed to the container for BELEAF 50SG insecticide.

CROP: GRAPES AND OTHER SMALL VINE CLIMBING FRUIT (EXCEPT FUZZY KIWIFRUIT) SUBGROUP 13-07F (7 DAY PHI)		
Includes Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and hybrids of these commodities.		
PESTS	RATE PER ACRE	USE DIRECTIONS
Aphids Plant Bugs (<i>Lygus spp.</i>)	2.8 to 4.2 oz (0.088 to 0.131 lb. a.i. / A)	Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines. Use LOWER RATE for building populations and use HIGHER RATE for greater populations and/or dense foliage. Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment. If identification of aphid species has not been confirmed, use HIGHER RATE.
Vine Mealybugs	4.2 to 5.6 oz (0.131 to 0.175 lb. a.i. / A)	Apply when mealybugs first appear, before populations increase to damaging levels. Application provides SUPPRESSION only. For control, apply in combination with other effective products labeled for use on these listed crops. Scout fields often and retreat as necessary to maintain populations below damaging levels.
SPRAY RECOMMENDATIONS Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 50 gallons per acre when applied by ground; use a minimum of 10 gallons per acres by air. Finished spray volumes should be increased under extreme pest populations or dense plant foliage. Spray adjuvants may improve coverage but do not use binder or sticker-type surfactants.		
USE RESTRICTIONS <ul style="list-style-type: none"> • Do not apply more than 1 application of Beleaf 50SG at the maximum single application rate of 5.6 oz/acre (0.175 lbs ai/acre) per year. • Do not apply more than 8.4 oz/acre of Beleaf 50SG (0.263 lbs. ai/acre) per year. This is equivalent to: <ul style="list-style-type: none"> o 3 applications at a low rate of 2.8 oz/acre Beleaf 50SG per year, or o 2 applications at a medium rate of 4.2 oz/acre Beleaf 50SG per year • Allow a minimum of 7 days between applications. 		

R-4729 062820 06-16-22

Manufactured for:



FMC Corporation
 2929 Walnut Street
 Philadelphia PA 19104

FMC – Trademark of FMC Corporation
 Beleaf® is a registered trademark of Ishihara Sangyo Kaisha, Ltd