

---

**SUPPLEMENTAL LABELING**

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1162A-S2 0920**

Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

<b>S-METOLACHLOR</b>	<b>GROUP</b>	<b>15</b>	<b>HERBICIDE</b>
<b>METRIBUZIN</b>	<b>GROUP</b>	<b>5</b>	<b>HERBICIDE</b>

**Boundary® 6.5EC Herbicide**

**This supplemental label expires on 09/15/2023 and must not be used or distributed after this date.**

Active Ingredients:

S-metolachlor*.....	58.2%
Metribuzin**.....	13.8%
<hr/>	
Other Ingredients***:	28.0%
Total:	100.0%

Boundary® 6.5EC Herbicide is formulated as an emulsifiable concentrate (EC) containing 5.25 lb of S-metolachlor and 1.25 lb of metribuzin per gallon.

\*CAS No. 87392-12-9

\*\*CAS No. 21087-64-9

\*\*\*Contains approximately 14% petroleum distillates.

**KEEP OUT OF REACH OF CHILDREN.**

## **WARNING/AVISO**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted in detalle. If you do not understand the label, find someone to explain it to you in detail.**

**EPA Reg. No. 100-1162**

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Boundary 6.5EC Herbicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**



## **DIRECTIONS FOR USE**

---

### **SOYBEANS (EXCEPT CALIFORNIA)**

---

Boundary 6.5EC Herbicide may be applied preplant surface, preplant incorporated, preemergence, postemergence directed or as a sequential application to control weeds listed on this label.

#### **Grazing and Feeding Treated Soybean Plants**

Treated soybean plants may be grazed or fed to livestock 40 days after the last application of Boundary 6.5EC Herbicide.

#### **Rate Ranges**

Where a rate range is shown, use a lower rate on soils that are coarse-textured and/or low in organic matter. Use a higher rate on soils that are relatively fine-textured and/or high in organic matter.

#### **Replanting**

If replanting is necessary in fields previously treated with Boundary 6.5EC Herbicide, the field may be replanted to soybeans. Excessive tillage may result in reduced weed control. Do not apply a second treatment as injury to soybeans may occur.

#### **Precautions (Soybeans)**

Injury to soybeans or reduced weed control may occur when Boundary 6.5EC Herbicide is used under the following conditions; avoid these conditions wherever possible.

- When soils have a calcareous surface area or a pH of 7.5 or higher.
- Due to the sensitivity of certain soybean varieties, there is an increased risk of crop injury with use of Boundary 6.5EC Herbicide on Altona, AP 55, AP 71, Asgrow 6520, Burlison, Coker 102, Coker 156, Dassel, GL 3202, Govan, Maple Amber, NB 3665, NKS 1884, Paloma 350, Portage, Regal, Semmes, Terra-Vig 505, Terra-Vig 606, Tracy, Vansoy, and Vinton 81. If you choose to plant a newly released soybean variety, consult your seed supplier for information on its tolerance to metribuzin (an active ingredient in Boundary 6.5EC Herbicide) before using Boundary 6.5EC Herbicide.
- When applied in conjunction with soil-applied organic phosphate pesticides.
- Uneven application or improper incorporation of Boundary 6.5EC Herbicide can decrease the level of weed control and/or increase the level of crop injury.
- When applied to any soil with less than 0.5% organic matter.
- Where soil incorporation is deeper than advised.

- When sprayers were not calibrated accurately.
- When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
- When soybeans are planted less than 1-½" deep, particularly when Boundary 6.5EC Herbicide is applied preemergence.
- Where high soil levels of atrazine are present.
- When using poor quality soybean seed.

**Restriction**

- Do not exceed 3.71 lb ai/A/year of S-metolachlor- containing products.

©2020 Syngenta