



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION

PROCURE® 480SC EPA Reg. No. 400-518

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF MICHIGAN, NEW JERSEY, PENNSYLVANIA, VIRGINIA, AND WEST VIRGINIA

This recommendation must be in the possession of the user at the time of application.

FOR CONTROL OF QUINCE RUST IN APPLES

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

DIRECTIONS FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. All applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA-registered label for Procure 480SC Fungicide (EPA Reg. No. 400-518) are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the Procure 480SC Fungicide label. This FIFRA Section 2(ee) recommendation must be in the possession of the user at the time of pesticide application.

CROP	DISEASE	RATE FL OZ PER ACRE (LBS AI/A)	APPLICATION INSTRUCTIONS
Apples	Scab (<i>Venturia inaequalis</i>) Cedar Apple Rust (<i>Gymnosporangium juniperi-virginianae</i>) Quince Rust (<i>Gymnosporangium clavipes</i>)	8 to 16 (0.25 to 0.50 lb ai)	Standard Spray Schedule: Initiate spraying at ½” green tip stage, continue through second cover or until primary scab sporulation is complete. Eradicant Spray Schedule (Post Infection Schedule): Initiate applications within 96 hours of the start of an infection period. Under low to moderate disease pressure conditions, rates in the lower range can be applied within 48 hours after the start of an infection period. Use rates in the higher range when disease pressure is high or if the time interval after the start of infection is greater than 48 hours. For improved scab control, apply PROCURE 480SC in combination with a protectant fungicide. After first cover, PROCURE 480SC can be used in a standard cover spray program in tank mixes with appropriate protectant fungicides.

RESTRICTIONS:

- Do not apply more than a total of 4 applications of PROCURE 480SC per crop per year.
- Do not apply more than a total of 56 fl oz (1.75 lb ai) of PROCURE 480SC per crop per year.
- Do not apply less than 14 days apart. Wait at least 21 days between second and third applications.
- Do not apply within 14 days of harvest.
- Do not allow livestock to graze on treated orchard ground cover.
- When making aerial applications, apply 12 – 16 fl oz PROCURE 480SC in a minimum of 10 gallons of water per acre.
- Do not apply by air in California.
- Ground applications – apply in a minimum of 100 gallons per acre.