



# Recommendation

## FIFRA SECTION 2(ee) RECOMMENDATION

### PROCURE<sup>®</sup> 480SC Fungicide EPA Reg. No. 400-518

#### FOR CONTROL OF BLACK ROT IN GRAPES FOR USE AND DISTRIBUTION IN THE STATES OF CONNECTICUT, DELAWARE, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, NEW HAMPSHIRE, NEW JERSEY, OHIO, PENNSYLVANIA, RHODE ISLAND, VERMONT, VIRGINIA, WEST VIRGINIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

SITE	DISEASE	RATE FL OZ PER ACRE (LBS AI/A)	APPLICATION INSTRUCTIONS
Grapes	Black Rot ( <i>Guignardia bidwellii</i> )	6 to 8 (0.19 to 0.50 lb ai)	Start protectant applications before bloom and continue on a 14 to 21-day interval. When disease pressure is high or for the most susceptible varieties, use 6 to 8 fl oz and/or 14-day spray intervals. When disease pressure is low or for least susceptible varieties, use 6 fl oz/acre and/or 21-day spray intervals.  Select spray volume, nozzle adjustments and ground speed to ensure complete coverage of fruit clusters and foliage. Use a minimum of 50 gallons of water per acre for early season applications.

#### RESTRICTIONS:

- Do not apply more than a total of 4 applications of PROCURE 480SC per crop per year.
- Do not apply more than a total of 32 fl oz (1.0 lb ai) of PROCURE 480SC per crop per year.
- Do not apply within 7 days of harvest.
- The Restricted Entry Interval (REI) for girdling table and raisin grapes is 24 hours (1-day).
- Do not allow by air.

Rev.3/4/2021