



A protectant fungicide for the control of powdery mildew in artichoke, gourds, lettuce, melons, pepper, pumpkin, stone fruits, strawberry, winter squash, cherry, grapes, and hops.

ACTIVE INGREDIENT:

quinoxyfen: 5,7-dichloro-4-(p-fluorophenoxy) quinoline:**22.58%**

OTHER INGREDIENTS:**77.42%**

TOTAL:**100.00%**

Contains 2.08 lb of active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

| FIRST AID | |
|---|---|
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| <p>Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-478-0798.</p> | |

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

NET CONTENTS:

EPA Reg. No. 33906-26-10163
EPA Est. No.



Distributed by:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not apply this product one-half hour before dusk to one-half hour after dawn in counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

PRODUCT INFORMATION

Quintec® fungicide is a protectant fungicide for the control of powdery mildew diseases. Quintec does not control existing or latent powdery mildew infections and must be applied on a protectant schedule (latency is the period between infection of plant tissues and the development of visible symptoms). Quintec is compatible with most commonly used insecticides, miticides and fungicides.

USE RESTRICTIONS

- Do not graze or harvest forage from cover crops within areas treated with Quintec.
- Do not aerially apply this product on stone fruits, cherry, grapes, or hops. Aerial application is permitted on other crops for which this product is registered for use.
- Chemigation: Do not apply this product through any type of irrigation system.
- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.
- Rainfast Requirement: When applied alone, Quintec is rainfast 1 hour after application.
- Not for use in greenhouses or nurseries.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

| Crops | Re-Planting Interval |
|--|----------------------|
| registered crops | no restrictions |
| all other crops grown for food or feed | 30 days |

MIXING DIRECTIONS

Quintec may be applied in tank mix combination with labeled rates of other pesticides, including fungicides, insecticides, or other chemicals used in the culture of the labeled crop provided: (1) the tank mix product is labeled for the timing and method of application for the intended use; and (2) tank mixing with Quintec or other pesticides is not prohibited by the label of the tank mix product.

Fill the spray tank to at least one-half the required spray volume and start agitation. Add (in order) dry flowable, wettable powder, and Quintec or other flowable (aqueous suspension) products to the spray tank. Then add the specified rate of emulsifiable concentrate and solution products. Allow time for thorough mixing after adding each product. Finish filling spray tank and maintain continuous agitation in the spray tank during mixing, filling and spraying operations.

Oil Sprays:

Oil sprays may be integrated as part of a powdery mildew management program. Consult local university extension for specific information.

Use of Surfactants:

A moderate rate of a registered nonionic or organosilicone surfactant may improve crop coverage. Penetrants and stickers are not recommended.

Tank Mixing Directions:

- Read carefully and follow all applicable use directions including application rates, use precautions, and limitations on the respective product labels.
- Do not exceed specified application rates or maximum use rates if products with the same active ingredient are applied together
- For tank mix partners packaged in water soluble packaging, do not use with products containing boron or mix in equipment previously used to apply boron-containing mixtures unless the tank and spray equipment have been adequately cleaned.
- Always perform a (jar) test to ensure compatibility of products used in tank mixtures.

Tank Mix Compatibility Testing:

To determine the physical compatibility of Quintec with other pesticides, perform a jar test. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination must not be used.

Disease Resistance Management

Cherry, Grapes, Hops, Stone Fruits:

Quintec is in mode of action Group 13. Do not make more than five applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec

Gourds, Lettuce, Melons, Pepper, Pumpkin, Strawberry, and Winter Squash: Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.

All Crops:

To delay development of fungicide/bactericide resistance:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Gowan Company representative.

APPLICATION DIRECTIONS

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, apply Quintec in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Cherry, Grape, Hops, Stone Fruits:

Ground: Apply uniformly by ground application in a minimum spray volume of 30 gallons per acre. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. Do not apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

Aerial: Do not aerially apply this product on cherry, grape, hops, and stone fruits.

Artichoke, Head and Leaf Lettuce, Melons, Pepper, Strawberry, Gourds and Pumpkin, Winter Squash:

Ground: For thorough spray coverage and optimum disease control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

Aerial: Apply in a minimum spray volume of 5 gallons of water per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Do not apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

Artichoke, Cherry, Grape, Head and Leaf Lettuce, Hops, Melons (Subgroup 9A), Pepper, Stone Fruits, Strawberry, Winter Squash, Gourds and Pumpkin

| Crop | Disease | Re-Treatment Interval (Days) | Rate of Quintec (fl oz/acre) ¹ | Quinoxifen rate (lb a.i./acre) |
|-----------------------------------|---|------------------------------|---|--------------------------------|
| artichoke | powdery mildew (<i>Leveillula taurica</i>) | 10 - 14 | 4 – 6 | 0.065 - 0.098 |
| cherry | powdery mildew (<i>Podosphaera clandestina</i>) | 10 - 14 | 7 | 0.114 |
| Grape | powdery mildew (<i>Erysiphe necator</i>) | 14 | 4 | 0.065 |
| | | 21 | 6.6 | 0.108 |
| head and leaf lettuce | powdery mildew (<i>Erysiphe cichoracearum</i>) | 10 – 14 | 4 – 6 | 0.065 - 0.098 |
| hops | powdery mildew (<i>Podosphaera macularis</i>) | 14 | 4 - 8.2 | 0.065 - 0.134 |
| melons (subgroup 9A) ² | powdery mildew (<i>Erysiphe cichoracearum</i> <i>Sphaerotheca fuliginea</i>) | 10 - 14 | 4 – 6 | 0.065 - 0.098 |
| pepper ³ | powdery mildew (<i>Leveillula taurica</i> , <i>Oidiopsis tuarica</i>) | 10 - 14 | 4 – 6 | 0.065 - 0.098 |
| stone fruits ⁴ | powdery mildew (<i>Sphaerotheca pannosa</i> , <i>Podosphaera leucotricha</i>) | 10 - 14 | 7 | 0.114 |
| Strawberry | powdery mildew (<i>Sphaerotheca macularis</i>) | 10 - 14 | 4 – 6 | 0.065 – 0.098 |
| winter squash, gourds, pumpkin | powdery mildew (<i>Erysiphe cichoracearum</i> <i>Sphaerotheca fuliginea</i>) | 10 - 14 | 4 – 6 | 0.065 -0.098 |

¹When applied as a dilute spray, maintain the per acre use rate.

²Melons (subgroup 9A) including citron melon, muskmelon (cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, true cantaloupe), watermelon

³Pepper including: bell pepper, chili pepper, cooking pepper, non-bell pepper, pimento, sweet pepper

⁴Stone fruits including apricot, chickasaw plum, damson plum, Japanese plum, nectarine, peach, plum, plumcot, prune (fresh)

SPECIFIC USE RESTRICTIONS

Artichoke

- **Preharvest Interval:** None
- **Maximum Application Rate:** Do not apply more than 32 fl oz of Quintec per acre per calendar year.
- Quintec is in mode of action Group 13. Do not make more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

Cherry

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or more than 32 fl oz of Quintec per acre per calendar year.
- **Minimum Spray Interval:** 7 days.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.

Grape

- **Preharvest Interval:** Do not apply within 21 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6.6 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.

- Quintec is in mode of action Group 13. Do not make more than five applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.

Head and Leaf Lettuce

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

Hops

- **Preharvest Interval:** Do not apply within 21 days of harvest.
- **Maximum Application Rate:** Do not apply more than 8.2 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.
- **Minimum Spray Interval:** 14 days.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.

Melons (Subgroup 9A)

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

Pepper

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per calendar year.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

Stone Fruits

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or 28 fl oz of Quintec per acre per calendar year.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

Strawberry

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after the first application of Quintec, discontinue its use.

Winter Squash, Gourds and Pumpkin

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 8 fl oz of Quintec per acre per application or 32 fl oz of Quintec per calendar year.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable containers 5 gallons or larger:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable containers 5 gallons or larger:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

FOR 24 HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.

For other product information, contact Gowan Company or see Safety Data Sheet.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Quintec® is a registered trademark of Nissan Chemical Corporation.

All other brands are registered trademarks of their respective owners.