

March 30, 2021

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc id: 572976

FIFRA 2(ee) Recommendation



**ADDING CONTROL OF ASCOCHYTA, BOTRYOSPHAERIA, MONILINIA, PHOMOPSIS, GUIGNARDIA,
SPUR BLIGHT and GUMMY STEM BLIGHT.
ADDING SUPPRESSION OF RUSTS OF SELECTED CROPS**

EPA REG. No. 91197-3-92488

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

Not for use in California or Texas

DIRECTIONS FOR USE

For control of Monilinia (*Monilinia* spp.) and Botryosphaeria (*Botryosphaeria* spp.) on berries and small fruits (e.g., blueberry (highbush and lowbush), blackberry, raspberry, cranberry)

For control of Phomopsis (*Phomopsis* spp.) on grapes, berries and small fruits (e.g., blueberry (highbush and lowbush), blackberry, raspberry, cranberry)

For control of Black rot (*Guignardia* spp.) on grapes

For control of Spur blight (*Didymella*) on raspberry

For control of Ascochyta blight (*Ascochyta* spp.) on beans, peas and dry pulses

For control of Gummy Stem Blight (*Stagonosporopsis* spp.) on cucurbit vegetables

For suppression of Rusts (*Puccinia* spp.) on small fruits

apply 5-7.5 lbs/A (80-120 oz/100 gal water) of Howler as a foliar spray or via chemigation. Use of an adjuvant is recommended. Apply in rotations or mixtures with other effective fungicides.

© 2021 AgBiome Innovations, Inc. All rights reserved.